

Brownian hitting distributions in space-time of bounded sets and the expected volume of Wiener sausage for Brownian bridges

Kôhei UCHIYAMA

Department of Mathematics, Tokyo Institute of Technology
Oh-okayama, Meguro Tokyo 152-8551
e-mail: uchiyama@math.titech.ac.jp

running head: Brownian hitting distributions in space-time

key words: Brownian hitting time, caloric measure, Wiener sausage, Brownian bridge.

AMS Subject classification (2010): Primary 60J65, Secondary 60J45. 35K20

Abstract

The space-time distribution, $Q_A(\mathbf{x}, dtd\xi)$ say, of Brownian hitting of a bounded Borel set A of $\mathbf{R}^d$ is studied. We derive the asymptotic form of the leading term of the time-derivative $Q_A(\mathbf{x}, dtd\xi)/dt$ for each $d = 2, 3, \dots$, valid uniformly with respect to the starting point $\mathbf{x}$ of the Brownian motion, which result significantly extends the classical ones for $Q_A(\mathbf{x}, dtd\xi)$ itself by Hunt ($d = 2$), Joffe and Spitzer ($d \geq 3$). The results obtained are applied to find the asymptotic form of the expected volume of Wiener sausage for the Brownian bridge joining the origin to a distant point.

1 Introduction and Main Results

In this paper we primarily focus on the space-time distribution of Brownian hitting of a bounded Borel set A of $\mathbf{R}^d$ expressed as

$$Q_A(\mathbf{x}, dtd\xi) = P_{\mathbf{x}}[B_{\sigma(A)} \in d\xi, \sigma_A \in dt] \quad (t > 0, d\xi \subset \partial A). \quad (1.1)$$

Here $P_{\mathbf{x}}$ denotes the law of a Brownian motion B_t started at $\mathbf{x}$, σ_A (or $\sigma(A)$) the first hitting time of A by B_t , namely $\sigma_A = \inf\{t > 0 : B_t \in A\}$, and ∂A the Euclidian boundary of A . When A is a disc ($d = 2$) or a ball ($d \geq 3$), the distribution $P_{\mathbf{x}}[\sigma_A < t]$ or its density $P_{\mathbf{x}}[\sigma_A \in dt]/dt$ are investigated by several recent works [8], [5], [23],[25] seeking the asymptotic behavior of them for $\mathbf{x} \notin A$ as $t \rightarrow \infty$. For general A the asymptotic form of the distribution $P_{\mathbf{x}}[\sigma_A < t]$ is given by Hunt [10] for $d = 2$ and by Joffe [13] and Spitzer [19] for $d \geq 3$. Their classical results may read as follows: if A is compact and $\mathbf{R}^d \setminus A$ is connected, then for each $\mathbf{x} \in \mathbf{R}^d \setminus A$, as $t \rightarrow \infty$

$$P_{\mathbf{x}}[t < \sigma_A] \sim \frac{2e_A(\mathbf{x})}{\lg t} \quad d = 2,$$

$$P_{\mathbf{x}}[t < \sigma_A < \infty] \sim \text{Cap}(A)P_{\mathbf{x}}[\sigma_A = \infty] \frac{t^{-d/2+1}}{(2\pi)^{d/2}(d/2-1)} \quad d \geq 3.$$

where e_A is a Green function for the open set $\mathbf{R}^d \setminus A$ with a pole at infinity and $\text{Cap}(\cdot)$ a Newtonian capacity (see (1.4) and (1.5) below). The latter result is uniform for $\mathbf{x}$ in each compact set of $\mathbf{R}^d \setminus A$, whereas for the former one, resting on a Tauberian theorem, uniformity

is not assured. In [10], [13] and [19] A is assumed compact as in above but the extension to bounded Borel sets is immediate with a knowledge about measurability question of the hitting time [11], [4], [1]. M. van den Berg [2] recently improves the result in the case $d \geq 3$ by obtaining sharp remainder estimates. In this paper we derive explicit asymptotic forms of the time derivative $Q_A(\mathbf{x}, dtd\xi)/dt$ as $t \rightarrow \infty$ valid uniformly for $|\mathbf{x}| = o(t)$ or for $t = o(|x|)$. By easy integration the asymptotic forms of $P_{\mathbf{x}}[t < \sigma_A < \infty]$ and $P_{\mathbf{x}}[\sigma_A < t | \sigma_A < \infty]$ can be computed from our results given below (see Section 6.5).

The measure kernel $Q_A(\mathbf{x}, dtd\xi)$ plays a significant role in the theory of heat operator. If A is compact and Ω_A denotes the unbounded component of $\mathbf{R}^d \setminus A$, then $Q_A(\mathbf{x}, dtd\xi)$ is identified with the lateral part of the caloric measure for the heat operator $\frac{1}{2}\Delta - \partial_t$ in the space-time domain $D = \{(\mathbf{x}, t) \in \mathbf{R}^d \times (0, \infty) : \mathbf{x} \in \Omega_A\}$, the exterior of a cylinder (see Section 6.1). (The other part of it is nothing but the measure whose density is given by the heat kernel for Ω_A with Dirichlet zero boundary condition.)

The present work is partly motivated and steered by a study of Wiener sausage swept by the set A attached to a d -dimensional Brownian motion started at the origin. Our interest is in finding a correct asymptotic form of the expected volume of the sausage of length t as $t \rightarrow \infty$ under the conditional law given that the Brownian motion at time t is at a given site $\mathbf{x}$ which is outside a parabolic region so that $\mathbf{x}^2 > \varepsilon t$ for any positive ε . To this end it is needed in our approach to estimate the density $Q_A(\mathbf{x}, dtd\xi)/dt$; we shall derive asymptotic forms of the expected volume by applying the results on Q_A . Fine estimates are obtained in the case when the process is pinned at the origin by McGillivray [15] ($d \geq 3$) and [16] ($d = 2$) (cf. also [3]) and in the case when $d = 2$, the value at time t is pinned within a parabolic region and A is a disc by [24]. There are many works for the sausage in the unconditional case (see, e.g., the references of [3]).

Let $U(a) \subset \mathbf{R}^d$ denote the open ball of radius $a > 0$ and centered at the origin. Let A be a bounded Borel set as above and denote by A^r the set of all regular points of A , i.e., those of $\mathbf{y} \in \bar{A}$ such that $P_{\mathbf{y}}[\sigma_A = 0] = 1$, and by Ω_A the unbounded component of $\mathbf{R}^d \setminus A^r$ in the fine topology, or what amounts to the same thing,

$$\Omega_A = \{\mathbf{x} \in \mathbf{R}^d : P_{\mathbf{x}}[\sigma_{\partial U(R)} < \sigma_A] > 0\}$$

for some (hence any) R such that $\bar{A} \subset U(R)$. (The over bar designates the Euclidean closure: $\bar{A} = A \cup \partial A$, where ∂A denotes the Euclidean boundary of A .) The set A^r is Borel and finely closed and $A \setminus A^r$ is polar, so that $P_{\mathbf{x}}[B_{\sigma(A)} \in A^r | \sigma_A < \infty] = 1$ for all $\mathbf{x}$ (see e.g. [17], [1]). Define for $\mathbf{x} \in \Omega_A$,

$$q_A(\mathbf{x}, t) = \frac{d}{dt} P_{\mathbf{x}}[\sigma_A \leq t]$$

and

$$H_A(\mathbf{x}, t; d\xi) = \frac{Q_A(\mathbf{x}, dtd\xi)}{dt} = \frac{P_{\mathbf{x}}[B_{\sigma(A)} \in d\xi, \sigma_A \in dt]}{dt} \quad (d\xi \subset \partial A, t > 0). \quad (1.2)$$

(Although both q_A and H_A depend on d , we do not designate this dependence in the notation.) Our purpose of this paper is to find the asymptotic form of $H_A(\mathbf{x}, t; \cdot)$ as $t \rightarrow \infty$. To this end it is often convenient to factor it into a product by conditioning on σ_A as follows:

$$H_A(\mathbf{x}, t; d\xi) = P_{\mathbf{x}}[B_{\sigma(A)} \in d\xi | \sigma_A = t] q_A(\mathbf{x}, t). \quad (1.3)$$

Put

$$p_t^{(d)}(x) = (2\pi t)^{-d/2} e^{-x^2/2t} \quad (x = |\mathbf{x}|)$$

and for $d \geq 3$

$$G^{(d)}(x) = \int_0^\infty p_t^{(d)}(x) dt = \frac{\Gamma(\nu)}{2\pi^{\nu+1} x^{2\nu}} \quad (\nu = \frac{1}{2}d - 1).$$

We usually write x for $|\mathbf{x}|$, $\mathbf{x} \in \mathbf{R}^d$ (as above) and sometimes $p_t^{(d)}(\mathbf{x})$ for $p_t^{(d)}(x)$ (and similarly for $G^{(d)}$) and $\sigma(A)$ for σ_A unless doing these causes any confusion. Denote by $\text{nb}_\varepsilon(A)$ the open ε -neighborhood of A in $\mathbf{R}^d$. We write $f(t) \sim g(t)$ if $f(t)/g(t) \rightarrow 1$ in any process of taking limit like ' $t \rightarrow \infty$ '.

The results of this paper are summarized in the following propositions (i) through (viii), where A is bounded Borel and non-polar, i.e., $P_{\mathbf{x}}[\sigma_A < \infty] > 0$ for $\mathbf{x} \notin \bar{A}$. We indicate in the square brackets at the head of each statement the theorem which the result presented below is taken from.

SUMMARY OF MAIN RESULTS.

I. Case $x/t \rightarrow 0$.

For each $\varepsilon > 0$, the following asymptotic formulae (i) and (ii) hold uniformly for $\mathbf{x} \in \Omega_A \setminus \text{nb}_\varepsilon(A^r)$:

(i) [Theorem 3.1] if $d \geq 3$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$,

$$q_A(\mathbf{x}, t) = \text{Cap}(A) P_{\mathbf{x}}[\sigma_A = \infty] p_t^{(d)}(x) (1 + o(1)), \quad (1.4)$$

where $\text{Cap}(A)$ denotes the Newtonian capacity of A (see Section 3.1.1) normalized so that

$$\text{Cap}(U(a)) = a^{d-2}/G^{(d)}(1);$$

(ii) [Theorem 3.2] if $d = 2$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$

$$q_A(\mathbf{x}, t) = p_t^{(2)}(x) \times \begin{cases} \frac{4\pi e_A(\mathbf{x})}{(\lg t)^2} \left(1 + O\left(\frac{1}{\lg t}\right)\right) & (x \leq \sqrt{t}), \\ \frac{\pi}{\lg(t/x)} \left(1 + O\left(\frac{1}{\lg(t/x)}\right)\right) & (x > \sqrt{t}), \end{cases} \quad (1.5)$$

where $e_A(\mathbf{x})$ denotes the Green function for Ω_A with a pole at infinity normalized so that

$$e_{U(a)}(\mathbf{x}) = \lg(x/a)$$

(cf. [14], p.369; see also Section 6.2));

(iii) [Theorem 3.3] for $d \geq 2$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$

$$\left| P_{\mathbf{x}}[B(\sigma_A) \in \cdot \mid \sigma_A = t] - H_A^\infty \right|_{\text{t.var}} \longrightarrow 0. \quad (1.6)$$

Here, H_A^∞ stands for the harmonic measure for the Brownian motion started at infinity conditioned on $\sigma_A < \infty$: $H_A^\infty(d\xi) = \lim_{x \rightarrow \infty} P_{\mathbf{x}}[B(\sigma_A) \in d\xi \mid \sigma_A < \infty]$; and $|\cdot|_{\text{t.var}}$ designates the total variation.

Let $S_A(t)$ denote the Wiener sausage of length t swept by A attached to a Brownian motion B_t :

$$S_A(t) = \{\mathbf{z} \in \mathbf{R}^d : \mathbf{z} - B_s \in A \text{ for some } s \in [0, t]\}.$$

The d -dimensional volume of $A \subset \mathbf{R}^d$ is denoted by $\text{vol}_d(A)$. Then

(iv) [Theorem 4.1] if $d \geq 3$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$

$$E_0 [\text{vol}_d(S_A(t)) \mid B_t = \mathbf{x}] \sim \text{Cap}(A)t; \quad (1.7)$$

(v) [Theorem 4.2] if $d = 2$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$,

$$E_0 [\text{vol}_2(S_A(t)) \mid B_t = \mathbf{x}] = \begin{cases} \frac{2\pi t}{\lg t}(1 + o(1)) & \text{if } |x| \leq \sqrt{t}, \\ \frac{\pi t}{\lg(t/x)}(1 + o(1)) & \text{if } |x| > \sqrt{t}. \end{cases} \quad (1.8)$$

II. Case $x/t \rightarrow v$.

In the case when x/t is bounded away from zero and infinity the results are not explicit as above. We can assert only the following one.

(vi) Let $d \geq 2$ and $v \geq 0$. Put $R_A = \sup\{|\mathbf{y}| : \mathbf{y} \in A^r\}$. There exists a measure kernel $\lambda_A(\mathbf{v}; d\xi)$ on $\mathbf{R}^d \times \partial A$ such that $\lambda_A(\mathbf{v}; \partial A)$ is positive, continuous and bounded (in $\mathbf{v}$) and such that as $x/t \rightarrow v > 0$ and $t \rightarrow \infty$,

$$\left| \frac{H_A(\mathbf{x}, t; \cdot)}{\Lambda_\nu(R_A x/t) p_t^{(d)}(x)} - \lambda_A(\mathbf{x}/t; \cdot) \right|_{\text{t.var}} \longrightarrow 0$$

and

$$E_0 [\text{vol}_d(S_A(t)) \mid B_t = \mathbf{x}] \sim t \Lambda_\nu(R_A x/t) \int_{\partial A} e^{-\xi \cdot \mathbf{x}/t} \lambda_A(\mathbf{x}/t; d\xi),$$

with $\Lambda_\nu(y)$ a positive function of y given in (2.1). Here, the convergence is uniform in $v \leq M$ in both the formulae for each $M > 1$ and they become continuously linked at $v = 0$ to the formulae of (iii) through (v). (See Section 4.2; λ_A is given in (4.14)). For the balls or discs the last formula may be given somewhat more explicitly:

$$E_0 [\text{vol}_d(S_{U(a)}(t)) \mid B_t = \mathbf{x}] \sim a^{2\nu} t \Lambda_\nu(av) \int_{\partial U(1)} e^{-a\xi \cdot \mathbf{x}/t} g_{av}(\theta) m_1(d\xi), \quad (1.9)$$

where $\theta \in [0, \pi]$ denotes the angle that $\xi \in \partial U(1)$ forms with $\mathbf{x}$ so that $\cos \theta = \xi \cdot \mathbf{x}/x$; and $g_\alpha(\theta)$, $\alpha \geq 0$ (given in (2.5) of Section 2) is a probability density relative to the uniform probability measure $m_1(d\xi)$ on $\partial U(1)$.

III. Case $x/t \rightarrow \infty$.

Let $\mathbf{e}$ be a unit vector of $\mathbf{R}^d$. Denote by $\Delta_{\mathbf{e}}$ the hyper-plane perpendicular to $\mathbf{e}$ passing through the origin and $\text{pr}_{\mathbf{e}} A$ the orthogonal projection of a set A on $\Delta_{\mathbf{e}}$. Let K be a compact set. Define a function $h = h_{\mathbf{e}, K}$ on $\Delta_{\mathbf{e}}$ by

$$h(\mathbf{z}) = \sup\{s \in \mathbf{R} : \mathbf{z} + s\mathbf{e} \in K\}, \quad \mathbf{z} \in \Delta_{\mathbf{e}}$$

with $\sup \emptyset = -\infty$ and let $\text{dis-ct}_{\mathbf{e}}(K)$ be the set of discontinuity points of h (see (3.36)). Suppose that

$$\text{vol}_{d-1}(\overline{\text{dis-ct}_{\mathbf{e}}(K)}) = 0,$$

where $\text{vol}_{d-1}(\cdot)$ denotes the $(d-1)$ -dimensional volume of a set of $\Delta_{\mathbf{e}}$. Then:

(vii) [Theorem 3.5] for $d \geq 2$, as $t \rightarrow \infty$ along with $v := x/t \rightarrow \infty$

$$\frac{H_K(x\mathbf{e}, t; d\xi)}{vp_t^{(d)}(x)e^{v\mathbf{e}\cdot\xi}} \Rightarrow m_{K,\mathbf{e}}(d\xi), \quad (1.10)$$

where ‘ $\Rightarrow$ ’ designates the weak convergence of measures and $m_{K,\mathbf{e}}$ stands for the Borel measure on ∂K induced from the $(d-1)$ -dimensional Lebesgue measure on $\text{pr}_{\mathbf{e}}K \subset \Delta_{\mathbf{e}}$ by the mapping $\mathbf{z} \in \text{pr}_{\mathbf{e}}K \mapsto \mathbf{z} + h(\mathbf{z})\mathbf{e} \in \partial K$ (see Section 3.2 for more details);

(viii) [Theorem 4.3] for $d \geq 2$, as $t \rightarrow \infty$ along with $x/t \rightarrow \infty$,

$$E_{\mathbf{0}}[\text{vol}_d(S_K(t)) \mid B_t = x\mathbf{e}] \sim \text{vol}_{d-1}(\text{pr}_{\mathbf{e}}K)x.$$

For convenience of later citation here we record the following scaling properties:

$$\begin{aligned} H_A(\mathbf{x}, t; d\xi) &= R^{-2}H_{R^{-1}A}(\mathbf{x}/R, t/R^2; R^{-1}d\xi) \quad (d\xi \subset \partial A), \\ q_A(\mathbf{x}, t) &= R^{-2}q_{R^{-1}A}(\mathbf{x}/R, t/R^2), \\ e_A(\mathbf{x}) &= e_{R^{-1}A}(\mathbf{x}/R) \quad (d=2) \quad \text{and} \\ \text{Cap}(A) &= R^{d-2}\text{Cap}(R^{-1}A) \quad (d \geq 3) \end{aligned} \quad (1.11)$$

($R > 0$). Note that the formula (1.4) is consistent to these relations.

The rest of the paper consists of five sections of the following contents.

§2. Hitting Distribution for A Disc/Ball.

§3. Hitting Distribution for A Bounded Borel Set.

3.1. Case $x/t \rightarrow 0$. 3.2. Case $x/\sqrt{t} \rightarrow \infty$.

§4. Expected Volume of Wiener Sausage for Brownian Bridge.

4.1. Case $x/t \rightarrow 0$. 4.2. Case $x/\sqrt{t} \rightarrow \infty$.

§5. Brownian Motion with Constant Drift $-v\mathbf{e}$.

§6. Appendix,

6.1. Harmonic measure of heat operator. 6.2. Uniform estimates for

$e_A(\mathbf{x})/P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]$. 6.3. An upper bound of q_A ($d \geq 3$). 6.4. Some

upper bounds of q_A ($d = 2$). 6.5. Asymptotics of the distribution of σ_A .

In §2 we summarize the results of [25] and [27] that are particularly relevant to the present subject. The results of both §3 and §4 heavily depend on those from §2.

The subjects of §3 and §4 are inter-related. Because of this the item (vi) is proved in Section 4.2. The asymptotic behavior of $H_A(\mathbf{x}, t; d\xi)$ is closely related to that of the expected volume of Wiener sausage swept by A attached to Brownian bridge joining $\mathbf{0}$ and $\mathbf{x}$. If $x/t \rightarrow 0$, the results on H_A entails those of the sausage. In the case $x/t \rightarrow \infty$, the situation is not so simple; our proof of (vii) relies on a result on the upper estimate of the expected volume of the sausage, and the lower estimate of it is obtained by using (vii).

Most of the statements advanced above may translate into the corresponding ones to Brownian motion with constant drift and some of them will be presented in §5. In §6 miscellaneous things are presented: a brief exposition of the well-known relation of the hitting distribution to caloric measure; some asymptotic evaluation of $P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]$ as $r \rightarrow \infty$ in terms of $e_A(\mathbf{x})$; some upper bounds of q_A ; asymptotic forms of $P_{\mathbf{x}}[\sigma_A \leq t]$.

2 Hitting Distribution for A Disc/Ball

Here we consider the case when $A = U(a)$, the open ball of radius a and centered at the origin, and state the asymptotic estimates of $H_{U(a)}(\mathbf{x}, t; d\xi)$ obtained in [25] and [27]. The following notation is used throughout the paper.

$$\begin{aligned} \nu &= \frac{d}{2} - 1 \quad (d = 1, 2, \dots); \\ \Lambda_\nu(y) &= \frac{(2\pi)^{\nu+1}}{2y^\nu K_\nu(y)} \quad (y > 0); \quad \Lambda_\nu(0) = \lim_{y \downarrow 0} \Lambda_\nu(y) \quad (\nu > 0). \end{aligned} \quad (2.1)$$

Here K_ν is the usual modified Bessel function (of second kind) of order ν . We write

$$q(x, t; a)$$

for $q_{U(a)}(\mathbf{x}, t)$. The definition of $q(x, t; a)$ may be naturally extended to Bessel processes of order ν and the results concerning it given below may be applied to such extension if $\nu \geq 0$. The following result from [25] provides a precise asymptotic form of the hitting time density for a ball.

Theorem 2.1. *Uniformly for $x > a$, as $t \rightarrow \infty$,*

$$q(x, t; a) = a^{2\nu} \Lambda_\nu\left(\frac{ax}{t}\right) p_t^{(d)}(x) \left[1 - \left(\frac{a}{x}\right)^{2\nu}\right] (1 + o(1)) \quad (d \geq 3) \quad (2.2)$$

and for $d = 2$,

$$q(x, t; a) = p_t^{(2)}(x) \times \begin{cases} \frac{4\pi \lg(x/a)}{(\lg(t/a^2))^2} (1 + o(1)) & (x \leq \sqrt{t}), \\ \Lambda_0\left(\frac{ax}{t}\right) (1 + o(1)) & (x > \sqrt{t}). \end{cases} \quad (2.3)$$

If the right-hand sides are multiplied by $e^{-a^2/2t}$, both the formulae (2.2) and (2.3) so modified hold true also as $x \rightarrow \infty$ uniformly for $t > 0$ (see also (2.4) below).

From known properties of $K_\nu(z)$ it follows that

$$\begin{aligned} \Lambda_\nu(y) &= (2\pi)^{\nu+1/2} y^{-\nu+1/2} e^y (1 + O(1/y)) \quad \text{as } y \rightarrow \infty; \\ \Lambda_\nu(0) &= 1/G^{(d)}(1) = 2\pi^{\nu+1}/\Gamma(\nu) = (d-2)\pi^{d/2}/\Gamma(d/2) \quad \text{for } \nu > 0; \text{ and} \\ \Lambda_\nu(y) &= \begin{cases} \frac{\pi}{-\lg(e^\gamma y/2)} (1 + O(y^2)) & (\nu = 0) \\ \Lambda_\nu(0) + O(y^2) & (\nu > 0, \nu \neq 1) \\ \Lambda_\nu(0) + O(y^2 \lg y) & (\nu = 1) \end{cases} \quad \text{as } y \downarrow 0. \end{aligned}$$

Here $\gamma = \int_0^1 (1 - e^{-t} - e^{-1/t}) t^{-1} dt$ (Euler's constant).

The following two results are also valid for Bessel processes of order $\nu \geq 0$. The first one is readily deduced from Theorem 2.1 by elementary computation (see the last section of [25]). The second one ([25] (Lemma 5), [5] (Lemma 4)) is also easily derived from the one-dimensional result with the help of a drift-transformation formula ((12) of [25]).

Theorem 2.2. For each $\nu \geq 0$, uniformly for $x > \sqrt{t/\lg t}$, as $t \rightarrow \infty$,

$$\left(\frac{a}{x}\right)^{-\nu} \int_0^t q(x, s; a) ds = \Lambda_\nu\left(\frac{ax}{t}\right) \frac{2^\nu}{(2\pi)^{\nu+1}} \int_{x^2/2t}^\infty e^{-y} y^{\nu-1} dy (1 + o(1)).$$

(Note that $(a/x)^\nu = \int_0^\infty q(x, s; a) ds$, so that the left side represents a conditional probability.)

Lemma 2.1. For each $\nu \geq 0$ it holds that uniformly for all $0 < t < a^2$ and $x > a$,

$$q(x, t; a) = \frac{x-a}{\sqrt{2\pi} t^{3/2}} e^{-(x-a)^2/2t} \left(\frac{a}{x}\right)^{(d-1)/2} \left[1 + O\left(\frac{t}{x}\right)\right]. \quad (2.4)$$

The case $x^2/t \rightarrow 0$ with $\lim(\lg x)/\lg t = 1/2$ of $\nu = 0$, not included in Theorem 2.2, is somewhat delicate. The following result is a reduced form of Theorem 3 of [23]).

Theorem 2.3. Let $\nu = 0$ and $\kappa = 2e^{-2\gamma}$. Uniformly for $\sqrt{t} > x > a$, as $t \rightarrow \infty$

$$\int_0^t q(x, s; a) ds = \frac{1}{\lg(\kappa t/a^2)} \left[1 - \frac{\gamma}{\lg(\kappa t/a^2)}\right] \int_{x^2/2t}^\infty \frac{e^{-y}}{y} dy + O\left(\frac{1}{(\lg t)^2}\right).$$

We sometimes need only the upper bound that follows immediately from Theorem 2.1 and it is convenient to write down it. In the following corollary, a result also obtained in [5], we include the bound for the case $0 < t < a^2$ and $d \geq 3$ that follows from Lemma 2.1.

Corollary 2.1. For all $a > 0, t > 0, x > a$,

$$q(x, t; a) \leq \kappa_d \left(1 - \frac{a}{x}\right) a^{2\nu} p_t^{(d)}(x-a) \left(1 \wedge \frac{t}{ax}\right)^{\nu-\frac{1}{2}}, \quad d \geq 3,$$

$$q(x, t; a) \leq \begin{cases} C \frac{\lg x/a}{t(\lg t/a^2)^2} & \text{for } x < \sqrt{t}, \\ C \frac{1}{1 + \lg(t/ax)} p_t^{(2)}(x) & \text{for } \sqrt{t} \leq x < t/a, \\ C \sqrt{\frac{ax}{t}} e^{ax/t} p_t^{(2)}(x) & \text{for } x \geq t/a \end{cases} \quad d = 2,$$

where κ_d is a constant depending only on d and C a universal constant. These bounds are optimal: the functions on the right-hand sides with some different positive constants provide lower bounds.

Next we present a result from [27] on the conditional distribution of the hitting site $B_{\sigma(U(a))}$ given $\sigma_{U(a)} = t$. Let $\theta = \theta(\xi) \in [0, \pi]$ denote the colatitude of a point $\xi \in \partial U(1)$ with $\mathbf{e}_0 = (1, 0, \dots, 0)$ taken to be the north pole, namely $\cos \theta = \xi \cdot \mathbf{e}_0$, and put

$$g_\alpha(\theta) = \sum_{n=0}^\infty \frac{K_0(\alpha)}{K_n(\alpha)} H_n(\theta). \quad (2.5)$$

for $\alpha > 0$ and $g_0(\theta) \equiv 1$. Here

$$H_n(\theta) = \begin{cases} \cos n\theta & \text{if } d = 2, \\ \kappa_{n,\nu} C_n^\nu(\cos \theta) & \text{if } d \geq 3, \end{cases}$$

where $\kappa_{n,\nu} = (n + \nu)\Gamma(\nu)/\sqrt{\pi}\Gamma(\nu + \frac{1}{2})$ and $C_n^\nu(z)$ is the Gegenbauer polynomial of order n associated with ν (cf.[29]). It follows that $H_0 \equiv 1$ and $g_\alpha(\theta)$ is jointly continuous in (α, θ) . Let $m_a(d\xi)$ denote the uniform probability measure on $\partial U(a)$.

Theorem 2.4. *Let $d \geq 2$. The function $g_\alpha(\theta)$ is positive on $[0, \pi]$, and for $v \geq 0$, as $x/t \rightarrow v$ and $t \rightarrow \infty$,*

$$\frac{P_{x\mathbf{e}_0}[B_{\sigma(U(a))} \in ad\xi \mid \sigma_{U(a)} = t]}{m_1(d\xi)} = \begin{cases} 1 + O(\ell_d(x, t)x/t) & \text{if } v = 0, \\ g_{av}(\theta)(1 + o(1)) & \text{if } v > 0 \end{cases} \quad (2.6)$$

uniformly for $(\xi, v) \in \partial U(1) \times [0, M]$ for each $M > 1$. Here $\ell_d(x, t) \equiv 1$ for $d \geq 3$ and

$$\ell_2(x, t) = (\lg t)^2 / \lg(2 + x) \quad (x < \sqrt{t}); \quad = \lg(t/x) \quad (x \geq \sqrt{t}).$$

Theorem 2.4 asserts that the limit distribution is uniform on the sphere $\partial U(a)$ if $\mathbf{x}/t \rightarrow 0$, while it is distributed with a positive and continuous density function $g_{av}(\theta)$ ($\theta = \theta(\xi)$) if $x/t \rightarrow v > 0$. In the case when x/t is unbounded, the situation becomes different: the weak limit concentrates at $a\mathbf{e}$ (see [27] for an exact asymptotic form of this case).

The next result, applied in Section 4.2.3, is a weak version of Theorem 2.4 of [27]; it is modified so as to be convenient for an application in this paper.

Theorem 2.5. *Let $d \geq 2$ and $t > 1$. Then as $v := x/t \rightarrow \infty$ and $\mathbf{x}/x \rightarrow \mathbf{e}$*

$$\frac{H_{U(a)}(\mathbf{x}, t; d\xi)}{\omega_{d-1} a^{2\nu} v p_t^{(d)}(x) e^{\xi \cdot \mathbf{x}/t}} \implies (\xi \cdot \mathbf{e}) m_a(d\xi) \mathbf{1}(\mathbf{e} \cdot \xi \geq 0), \quad (2.7)$$

where $\mathbf{1}(\mathcal{S})$ is 1 or 0 according as a statement $\mathcal{S}$ is true or false; the arrow ' $\implies$ ' signifies the weak convergence of finite measures (on $\partial U(a)$); and ω_n stands for the area of n -dimensional unit sphere.

It is noted that the convergence in (2.7) is necessarily uniform for $\mathbf{e} \in \partial U(a)$.

3 Asymptotics of H_A for A Bounded Borel Set

Here we seek an exact asymptotic form, as $t \rightarrow \infty$, of $H_A(\mathbf{x}, t; d\xi)dt$ for $\mathbf{x} \in \Omega_A$, where A is a bounded Borel set of $\mathbf{R}^d$.

3.1 Case $x/t \rightarrow 0$

We deal with cases $d \geq 3$ and $d = 2$ separately. In the case $d \geq 3$ the capacity of A is involved in the leading term, while for $d = 2$, the logarithmic capacity appears only in the next order term that we shall not identify.

Our basic strategy is to use the Huygens property of H_A in the form

$$H_A(\mathbf{x}, t; E) = \int_0^t ds \int_{\partial U(R)} H_{U(R)}(\mathbf{x}, t - s; d\xi) H_A(\xi, s; E) \quad (3.1)$$

for $E \subset \bar{A}$, $\mathbf{x} \notin U(R)$

valid if $A \subset U(R)$. For the Brownian motion started at a point very distant from A to hit A for the first time at time t it must hit $U(R)$ (with $R/x \ll 1$) for the first time in a relatively small time interval just before t with high probability, so that the outer integral of (3.1) must concentrate on such an interval and the exact asymptotic forms for balls $U(R)$ described in

the preceding section will yield the results for general A . In the case when the starting point is not distant from A the process must make a big excursion (in the interval $[0, \sigma_A]$) and exit from a large sphere within a relatively short time interval so that the problem is reduced to the case of distant starting points.

3.1.1. DENSITY OF HITTING TIME DISTRIBUTION ($d \geq 3$). Let $d = 3, 4, \dots$ and $\text{Cap}(A)$ denote the Newtonian capacity of A . For the present purpose it is convenient to define it by

$$\text{Cap}(A) = \frac{1}{G^{(d)}(1)} \lim_{x \rightarrow \infty} x^{2\nu} P_{\mathbf{x}}[\sigma_A < \infty].$$

Put

$$R_A = \sup\{|\mathbf{y}| : \mathbf{y} \in A\}.$$

Then the identity

$$x^{2\nu} P_{\mathbf{x}}[\sigma_A < \infty] = R^{2\nu} E_{\mathbf{x}}[P_{B_{\sigma(U(R))}}[\sigma_A < \infty] \mid \sigma_{U(R)} < \infty]$$

holds true whenever $x > R \geq R_A$ and shows the existence of the limit defining $\text{Cap}(A)$ as well as the formula

$$\text{Cap}(A) = \text{Cap}(U(R)) P_{m_R}[\sigma_A < \infty], \quad (3.2)$$

where m_R denotes the uniform probability measure on the sphere $\partial U(R)$. (See also Remark 2 (d) below.) We designate by c_R, c'_R etc. constants depending only on R_A and d whose exact values are not significant for the present purpose and may vary at different occurrences of them.

Theorem 3.1. *Let $d \geq 3$. Uniformly for $\mathbf{x} \in \Omega_A$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$*

$$q_A(\mathbf{x}, t) = \text{Cap}(A) \left(P_{\mathbf{x}}[\sigma_A = \infty] p_t^{(d)}(x) (1 + o(1)) + \text{err}(\mathbf{x}, t) \right), \quad (3.3)$$

where if $x \geq 2R_A$, $\text{err}(\mathbf{x}, t) = 0$ and if $x < 2R_A$,

$$|\text{err}(\mathbf{x}, t)| \leq \frac{c_R}{t^{d/2}} P_{\mathbf{x}}[\sigma_{A \cup \partial U(2R_A)} > t/\lg t] \leq c'_R e^{-\lambda t/\lg t} \quad (3.4)$$

with some $\lambda > 0$; $o(1)$ and λ as well as c_R, c'_R depend only on R_A and d . (The exponent $-\lambda t/\lg t$ in the upper bound for $\text{err}(\mathbf{x}, t)$ may be replaced any function that is $o(t)$.)

REMARK 2. (a) In view of our definition of $\text{Cap}(A)$ and Theorem 2.1 the formula (3.3) entails that as $x \rightarrow \infty$ and $x/t \rightarrow 0$

$$q_A(\mathbf{x}, t) = P_{\mathbf{x}}[\sigma_A < \infty] x^{2\nu} q(x, t; 1) (1 + o(1)),$$

which, on noting that $1/x^{2\nu} = P_{\mathbf{x}}[\sigma_{U(1)} < \infty]$, may be paraphrased in terms of conditional probability as follows: as $x \rightarrow \infty$ and $x/t \rightarrow 0$

$$P_{\mathbf{x}}[\sigma_A \in dt \mid \sigma_A < \infty]/dt = P_{\mathbf{x}}[\sigma_{U(1)} \in dt \mid \sigma_{U(1)} < \infty]/dt (1 + o(1)).$$

This formula does not hold in general if x/t is bounded away from zero.

(b) As $\mathbf{x}$ gets close to A and t large $q_A(\mathbf{x}, t)$ itself approaches zero very fast, but it is not so simple a matter to express in general how fast it does. In any case, in (3.3) the error

term $\text{err}(\mathbf{x}, t)$, though very small for t large (see (3.4)), cannot be absorbed into $o(1)$ that precedes it since for each $t > 1$, $P_{\mathbf{x}}[\sigma_{A \cup \partial U(2R_A)} > t]$ may be much larger than $P_{\mathbf{x}}[\sigma_A = \infty]$ (for some $\mathbf{x}$). An example is easily constructed by considering a ball pockmarked by infinitely many cave-like holes, each one containing a relatively spacial chamber connected by a narrow tunnel to the outside and among them there being one such that the ratio of diameter of the tunnel to that of the chamber is smaller than any prescribed number. For each $\varepsilon > 0$, $\inf_{\mathbf{x} \in \Omega_A \setminus \text{nb}\delta_\varepsilon(A^r)} P_{\mathbf{x}}[\sigma_A = \infty] > 0$, and hence the asymptotic formula

$$q_A(\mathbf{x}, t) = \text{Cap}(A) P_{\mathbf{x}}[\sigma_A = \infty] p_t^{(d)}(x) (1 + o(1)) \quad (3.5)$$

holds uniformly for $\mathbf{x} \in \Omega_A \setminus \text{nb}\delta_\varepsilon(A^r)$. With this remark one may readily verify that (3.5) holds uniformly for $\mathbf{x} \in \Omega_A$ if A satisfies some regularity condition such as smoothness of boundary.

(c) We can find a constant $\eta > 0$ such that under the constraint $x < \sqrt{2t \lg t}$, the factor $(1 + o(1))$ may be replaced by $(1 + O(t^{-\eta}))$ in (3.3). A sketch of proof is given at the end of this subsection.

(d) If μ_A is the equilibrium charge of A , namely the measure on $\bar{A}$ such that $P_{\mathbf{x}}[\sigma_A < \infty] = \int G^{(d)}(\mathbf{y} - \mathbf{x}) \mu_A(d\mathbf{y})$, then the capacity of A is usually defined as the total charge of μ_A . Our definition conforms to it as is well known:

$$\text{Cap}(A) = R^{2\nu} \int m_R(d\xi) \int G^{(d)}(\mathbf{y} - \xi) \mu_A(d\mathbf{y}) = \mu_A(\bar{A}).$$

A substantial part of the proof of Theorem 3.1 is contained in that of the next lemma.

Lemma 3.1. *As $x \rightarrow \infty$ and $x/t \rightarrow 0$*

$$q_A(\mathbf{x}, t) = \frac{\text{Cap}(A)}{\text{Cap}(U(1))} q(x, t; 1) (1 + o(1)). \quad (3.6)$$

Here $o(1)$ depends only on R_A and d .

Proof. Put $R = 2R_A$. Suppose $x \rightarrow \infty$. By strong Markov property

$$q_A(\mathbf{x}, t) = \int_0^t ds \int_{\partial U(R)} H_{U(R)}(\mathbf{x}, t - s; d\xi) q_A(\xi, s).$$

Take a (large) number M from the interval $(1, \sqrt{t})$ and split the range of outer integral at $s = M$ and $s = t/2$. We write

$$I_{[0, M]}, \quad I_{[M, t/2]} \quad \text{and} \quad I_{[t/2, t]}$$

for the corresponding integrals over the Cartesian products $[0, M] \times \partial U(R)$, etc. On employing Theorem 2.4 (with $v = 0$),

$$I_{[0, M]} = \int_0^M q(x, t - s; R) ds \int_{\partial U(R)} q_A(\xi, s) m_R(d\xi) \left(1 + O\left(\frac{x}{t}\right)\right). \quad (3.7)$$

Noting $p_{t-s}^{(d)}(x)/p_t^{(d)}(x) = \exp\{-\frac{x^2 s}{2t(t-s)}\} (1 + o(1))$ we apply Theorem 2.1 to see

$$\frac{q(x, t - s; R)}{q(x, t; R)} = \exp\left\{-\frac{x^2 s}{2t(t-s)}\right\} (1 + o(1)) \quad (3.8)$$

as $s/t \rightarrow 0$, whereas

$$\int_0^M ds \int_{\partial U(R)} q_A(\xi, s) m_R(d\xi) = P_{m_R}[\sigma_A < M] = \frac{R^{-2\nu} \text{Cap}(A)}{\text{Cap}(U(1))} (1 + O(M^{-\nu}))$$

in view of (3.2) (see Lemma 6.2 for the second equality). Substitute these relations into (3.7) and let $M \rightarrow \infty$ under $(1 \vee x^2)M/t^2 \rightarrow 0$. Then, observing

$$R^{-2\nu} q(x, t; R) = q(x, t; 1)(1 + o(1))$$

we find that $I_{[0, M]}$ is asymptotic to the right-hand side of (3.6) (provided $x/t \rightarrow 0$).

The integral over $[M, t/2] \times \partial U(R)$ is estimated as

$$I_{[M, t/2]} \leq \left(\sup_{M < s < t/2} \frac{q(x, t-s; R)}{q(x, t; R)} \sup_{\xi \in \partial U(R)} P_\xi[M < \sigma_A < \infty] \right) q(x, t; R),$$

hence negligible, for the first supremum is bounded and the second one is $\text{Cap}(A) \times O(M^{-\nu})$, hence negligible as $M \rightarrow \infty$.

It remains to ascertain that $I_{[t/2, t]}$ is also negligible. We prove that

$$I_{[t/2, t]} \leq \text{Cap}(A) q(x, t; R) \times c_R x^{-2\nu} e^{-2x^2/3t} \quad (R < x < t/R_A). \quad (3.9)$$

The proof rests on the bound

$$\sup_{s \geq t/2} q_A(\mathbf{y}, s) \leq \kappa_d \text{Cap}(A) t^{-d/2} \quad (3.10)$$

valid for all $t > R^2$ and $\mathbf{y} \in \Omega_A$, where the constant κ_d depends on d only. This bound plays a key role also later (see (3.18)), but its proof is somewhat involved and postponed to the last section (see Lemma 6.6). Clearly we have

$$I_{[t/2, t]} \leq P_{\mathbf{x}}[\sigma_{U(R)} < t/2] \sup_{t/2 \leq s \leq t} \sup_{|\xi|=R} q_A(\xi, s).$$

On taking (3.10) for granted and employing Theorem 2.2 the right-hand side of this inequality is dominated by

$$\begin{aligned} & \frac{\kappa_d \text{Cap}(A)}{t^{d/2}} \times c'_R P_{\mathbf{x}}[\sigma_{U(R)} < \infty] \int_{x^2/t}^{\infty} e^{-u} u^{\nu-1} du \\ & \leq c''_R \text{Cap}(A) \frac{[e^{-x^2/t} (x^2/t)^{\nu-1}] \wedge 1}{t^{d/2} x^{2\nu}}, \end{aligned} \quad (3.11)$$

which verifies (3.9) as required. The proof of Lemma 3.1 is complete. $\square$

Let $R = 2R_A$. For $\mathbf{x} \in \Omega_A \cap U(R)$, define $t_* = t_*(\mathbf{x}; A)$ via

$$P_{\mathbf{x}}[\sigma_{A \cup \partial U(R)} > t_*] = \frac{1}{2} P_{\mathbf{x}}[\sigma_A = \infty]. \quad (3.12)$$

Lemma 3.2. *Let $R, \mathbf{x}$ and t_* be as above. Then for $t > t_*$,*

- (i) $P_{\mathbf{x}}[\sigma_{\partial U(R)} \leq t \mid \sigma_A = \infty] \geq 1/2$; and
- (ii) $P_{\mathbf{x}}[\sigma_{\partial U(R)} > t \mid \sigma_A = \infty] \leq C e^{-\lambda(t-t_*)/R^2}$,

where C and λ are universal positive constants.

Proof. From the defining relation (3.12) it follows that

$$\frac{1}{2} = \frac{P_{\mathbf{x}}[\sigma_{A \cup \partial U(R)} > t_*]}{P_{\mathbf{x}}[\sigma_A = \infty]} \geq P_{\mathbf{x}}[\sigma_{\partial U(R)} > t_* \mid \sigma_A = \infty].$$

Hence $P_{\mathbf{x}}[\sigma_{\partial U(R)} \leq t_* \mid \sigma_A = \infty] \geq 1/2$, showing (i).

Put $u(\mathbf{y}, t) = P_{\mathbf{y}}[\sigma_{\partial U(R)} > t, \sigma_A = \infty]$. Then for $t > t_*$,

$$\begin{aligned} u(\mathbf{x}, t) &= E_{\mathbf{x}}[u(B_{t_*}, t - t_*); \sigma_{A \cup \partial U(R)} > t_*] \\ &\leq \frac{1}{2} P_{\mathbf{x}}[\sigma_A = \infty] \sup_{\mathbf{y} \in U(R)} u(\mathbf{y}, t - t_*) \\ &\leq C e^{-\lambda(t-t_*)/R^2} P_{\mathbf{x}}[\sigma_A = \infty]. \end{aligned}$$

Thus (ii) is obtained. $\square$

Proof of Theorem 3.1. In view of Lemma 3.1 (together with $\text{Cap}(U(1)) = \Lambda_{\nu}(0)$) and Theorem 2.1 we may suppose that x remains in a bounded set as $t \rightarrow \infty$ since $\lim_{x \rightarrow \infty} P_{\mathbf{x}}[\sigma_A = \infty] = 1$. The proof is based on

$$\begin{aligned} q_A(\mathbf{x}, t) &= \int_0^{\frac{3}{4}t} \int_{\partial U(r)} P_{\mathbf{x}}[\sigma_{\partial U(r)} \in ds, B_{\sigma(\partial U(r))} \in d\xi, \sigma_A > s] q_A(\xi, t - s) \\ &\quad + P_{\mathbf{x}}[\sigma_{\partial U(r)} > \frac{3}{4}t, \sigma_A \in dt]/dt, \end{aligned} \quad (3.13)$$

in which we let $r \rightarrow \infty$ under the constraint $r^2/t \rightarrow 0$ (so that $p_t^{(d)}(x) \sim p_t^{(d)}(r) \sim (2\pi t)^{-d/2}$). The first term on the right-hand side above contains the principle part and is evaluated in the step 1, while the second term is negligible in comparison to the first and dealt with in Step 2.

Step 1. First we consider the repeated integral above. It follows from Lemma 3.1 that uniformly for $\xi \in \partial U(r)$ and $s = o(t)$, as $r \rightarrow \infty$ in a way specified above

$$q_A(\xi, t - s) = \frac{\text{Cap}(A)}{\text{Cap}(U(1))} q(x, t; 1)(1 + o(1)). \quad (3.14)$$

On the other hand defining $\eta(\mathbf{x}, t, T)$ via

$$\begin{aligned} \int_0^T P_{\mathbf{x}}[\sigma_{\partial U(r)} \in ds, \sigma_A > s] &= P_{\mathbf{x}}[\sigma_{\partial U(r)} < T \wedge \sigma_A] \\ &= P_{\mathbf{x}}[\sigma_A = \infty] - \eta(\mathbf{x}, t, T), \end{aligned}$$

we observe that for every $T > 1$

$$-P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A < \infty] \leq \eta(\mathbf{x}, t, T) \leq P_{\mathbf{x}}[\sigma_{\partial U(r)} \geq T, \sigma_A = \infty], \quad (3.15)$$

in which the left-most member divided by $P_{\mathbf{x}}[\sigma_A = \infty]$ tends to zero as $r \rightarrow \infty$. We let $T \rightarrow \infty$ in such a manner that $r^2/T \rightarrow 0$ and $T/t \rightarrow 0$, the latter ensuring that (3.14) is valid for $s < T$. We need to deal with the cases $x \geq R$ and $x < R$ separately.

Let $x \geq R$. Noting $\inf_{|\mathbf{y}|=R} P_{\mathbf{y}}[\sigma_A = \infty] \geq 2^{-2\nu}$, and $P_{\mathbf{x}}[\sigma_{\partial U(r)} \geq T] = O(e^{-\lambda T/r^2})$ (for some $\lambda > 0$), one observes that

$$P_{\mathbf{x}}[\sigma_{\partial U(r)} > T \mid \sigma_A = \infty] = O(e^{-\lambda T/r^2}) \quad (3.16)$$

and then that

$$\frac{\eta(\mathbf{x}, t, T)}{P_{\mathbf{x}}[\sigma_A = \infty]} \rightarrow 0 \quad (3.17)$$

locally uniformly for $\mathbf{x} \in \Omega_A \setminus U(R)$. Hence, if we take e.g. $r = t^{1/4}$ and let $T \rightarrow \infty$ so that both r^2/T and T/t approach 0, then the integral in (3.13) over $[0, T]$ may be equated with the formula on the right-hand side of (3.3) with the term $\text{err}(\mathbf{x}, t)$ being discarded.

That the integral over $[T, 3t/4]$ is negligible is verified by noting that

$$\frac{P_{\mathbf{x}}[T \leq \sigma_{\partial U(r)} < \sigma_A]}{P_{\mathbf{x}}[\sigma_A = \infty]} \leq \frac{2P_{\mathbf{x}}[\sigma_{\partial U(r)} > T, \sigma_A = \infty]}{P_{\mathbf{x}}[\sigma_A = \infty]} = O(e^{-\lambda T/r^2}) \quad (3.18)$$

(in view of (3.16)) and then by applying (3.10) with $|\mathbf{y}| = r$.

Let $\mathbf{x} \in \Omega_A \cap U(R)$. In this case the obvious reasoning for (3.16) (hence (3.17)) breaks down. Let t_* be defined by (3.12). Then, if $T/2 \leq t_*$,

$$P_{\mathbf{x}}[\sigma_A = \infty] \leq 2P_{\mathbf{x}}[\sigma_{A \cup \partial U(R)} > T/2] = O(e^{-\lambda T/2R^2}). \quad (3.19)$$

On the other hand, if $T/2 > t_*$, by virtue of Lemma 3.2 we see that

$$\begin{aligned} & P_{\mathbf{x}}[\sigma_{\partial U(r)} > T, \sigma_A = \infty] \\ &= P_{\mathbf{x}}[\sigma_{\partial U(R)} \leq T/2, T < \sigma_{\partial U(r)} < \sigma_A] + P_{\mathbf{x}}[\sigma_{\partial U(R)} > T/2, \sigma_A = \infty] \\ &\leq P_{\mathbf{x}}[\sigma_{\partial U(R)} < \sigma_A] \sup_{\xi \in \partial U(R)} P_{\xi}[\sigma_{\partial U(r)} > T/2] + P_{\mathbf{x}}[\sigma_{A \cup \partial U(R)} > T/2] \\ &\leq CP_{\mathbf{x}}[\sigma_A = \infty]e^{-\frac{1}{2}\lambda T/r^2} + P_{\mathbf{x}}[\sigma_{A \cup \partial U(R)} > T/2]. \end{aligned}$$

Together with (3.19) this yields, instead of (3.17), the bound

$$|\eta(\mathbf{x}, t, T)| \leq CP_{\mathbf{x}}[\sigma_A = \infty]e^{-\frac{1}{2}\lambda T/r^2} + 3P_{\mathbf{x}}[\sigma_{A \cup \partial U(R)} > T/2]. \quad (3.20)$$

In view of the inequality in (3.18) the corresponding bound for the integral over $[T, 3t/4]$ is also valid and, recalling (3.14), we find the required estimate of the repeated integral in (3.13) being obtained by taking $T = 2t/\lg t$ and $r = t^{1/4}$: the second term on the right side of (3.20) is to turn out to be the main cause for the error estimate in (3.4).

Step 2. It remains to evaluate $P_{\mathbf{x}}[\sigma_{\partial U(r)} > \frac{3}{4}t, \sigma_A \in dt]/dt$, the second term on the right-hand side of (3.13). We decompose $\{\sigma_{\partial U(r)} > \frac{3}{4}t\} = \{\sigma_{\partial U(r)} > \frac{3}{4}t, \sigma_{\partial U(R)} > \frac{1}{2}t\} + \{\sigma_{\partial U(r)} > \frac{3}{4}t, \sigma_{\partial U(R)} \leq \frac{1}{2}t\}$ with $R = 2R_A$. Let I and II be the contributions of the first and the second ones of the decomposition, respectively. We first deal with II , which may be expressed as

$$\begin{aligned} & \frac{P_{\mathbf{x}}[\sigma_{\partial U(r)} > \frac{3}{4}t, \sigma_{\partial U(R)} \leq \frac{1}{2}t, \sigma_A \in dt]}{dt} \\ &= \int_0^{t/2} ds \int_{\partial U(R)} H_{A \cup \partial U(R)}(\mathbf{x}, s; d\xi) \\ & \quad \times \int_{U(r)} P_{\xi}[B_{\frac{1}{2}t-s} \in d\mathbf{y}, \sigma_{A \cup \partial U(r)} > \frac{3}{4}t - s] q_A(\mathbf{y}, \frac{1}{4}t + s). \end{aligned}$$

On using (3.10) and noting that

$$\int_0^{t/2} H_{A \cup \partial U(R)}(\mathbf{x}, s; \partial U(R)) ds \leq P_{\mathbf{x}}[\sigma_{\partial U(R)} < \sigma_A] \leq 2^{2\nu} P_{\mathbf{x}}[\sigma_A = \infty],$$

it is readily inferred that $II \leq \text{Cap}(A)P_{\mathbf{x}}[\sigma_A = \infty] \times o(1/t^{d/2})$.

As for the contribution of the first event which is contained in $\{\sigma_{\partial U(R)} > \frac{1}{2}t\}$, we observe that if $x < R$,

$$\begin{aligned} I &\leq \frac{P_{\mathbf{x}}[\sigma_{\partial U(R)} > t/2, \sigma_A \in dt]}{dt} \\ &= \int_{U(R)} P_{\mathbf{x}}[B_{t/2} \in d\mathbf{y}, \sigma_{A \cup \partial U(R)} > t/2] q_A(\mathbf{y}, t/2) \\ &\leq c_R \text{Cap}(A) P_{\mathbf{x}}[\sigma_{A \cup \partial U(R)} > t/2] t^{-d/2}, \end{aligned}$$

where the last inequality follows from (3.10). This may be absorbed into the error term in (3.3). If $x \geq R$,

$$\begin{aligned} I &\leq P_{\mathbf{x}}[\sigma_{U(R)} \wedge \sigma_{\partial U(r)} > t/2] \sup_{R \leq |\mathbf{y}| \leq r} q_A(\mathbf{y}, t/2) \\ &\leq C P_{\mathbf{x}}[\sigma = \infty] \text{Cap}(A) e^{-\lambda t/r^2}, \end{aligned}$$

where we have applied the bound $q_A(\mathbf{x}, t) \leq \kappa_d \text{Cap}(A)$ ($x > R$) (see Lemma 6.3 (ii)). Thus the required bound is obtained.

The proof of Theorem 3.1 is now complete. $\square$

Sketch of the proof of Remark 2 (c). Let $d \geq 3$. Theorem 3 of [25] provides an error estimate for the asymptotic form of $q(x, t; a)$ given in Theorem 2.1 and according to it the error term $o(1)$ in (2.2) can be replaced by $O(\lg t/t)$ for $x < \sqrt{2t \lg t}$ (the asymptotic form being exact if $d = 3$). On examining the proofs of Lemma 3.1 this shows that under the same constraint on x , (3.6)) can be refined to

$$q_A(\mathbf{x}, t) = \text{Cap}(A) p_t^{(d)}(x) (1 + O(x^{-2\nu} \vee t^{-\eta})) \quad (x < \sqrt{2t \lg t}) \quad (3.21)$$

hence $o(1)$ in (3.14) to $O(r^{-2\nu} \vee t^{-\eta})$, for some $\eta > 0$. The rest is done by noting that $P_{\mathbf{x}}[\sigma_A = \infty] = 1 + O(x^{-2\nu})$, the argument in the proof of Theorem 3.1 is valid also for e.g. $R < x < t^{1/5}$, and the terms appearing in the form $e^{-\lambda T/r^2}$ therein are all negligible.

3.1.2. DENSITY OF HITTING TIME DISTRIBUTION ($d = 2$). Let A be a bounded Borel set of $\mathbf{R}^2$ that is non-polar. Define

$$e_A(\mathbf{x}) = \pi \lim_{|\mathbf{y}| \rightarrow \infty} g_{\Omega_A}(\mathbf{x}, \mathbf{y}), \quad (3.22)$$

where $g_{\Omega_A}(\mathbf{x}, \mathbf{y})$ denotes the Green function for Ω_A , which is continuous on $\Omega_A \times \Omega_A$ (see Section 6.2). If A contains a curve connecting the origin with $\partial U(R_A)$, then $0 \leq e_A(\mathbf{x}) \leq C_0$ if $|\mathbf{x}| \leq R_A$, with a universal constant C_0 as being assured by the scaling property for e_A ; otherwise $\sup_{\mathbf{x} \in U(R_A)} e_A(\mathbf{x})$ could be arbitrarily large (depending on A). Further properties of e_A are reviewed in Section 6.2.

Theorem 3.2. *Let $d = 2$ and suppose A to be non-polar. Then, as $x \rightarrow \infty$ and $x/t \rightarrow 0$*

$$q_A(\mathbf{x}, t) = p_t^{(2)}(x) \times \begin{cases} \frac{4\pi \lg x}{(\lg t)^2} (1 + o(1)) & (x \leq \sqrt{t}), \\ \frac{\pi}{\lg(t/x)} \left(1 + O\left(\frac{1}{\lg(t/x)}\right)\right) & (x > \sqrt{t}). \end{cases} \quad (3.23)$$

Moreover, uniformly in $\mathbf{x} \in \Omega_A \cap U(\sqrt{t})$ as $t \rightarrow \infty$

$$q_A(\mathbf{x}, t) = \frac{4\pi e_A(\mathbf{x})}{(\lg t)^2} p_t^{(2)}(x) \left(1 + O\left(\frac{1}{\lg t}\right)\right) + \text{err}(\mathbf{x}, t) \quad (3.24)$$

with

$$|\text{err}(\mathbf{x}, t)| \leq \frac{c_R}{t(\lg t)^2} P_{\mathbf{x}}[\sigma_{A \cup \partial U(2R_A)} > t/2] \leq c'_R e^{-\lambda t} \quad (x < 2R_A)$$

for some $\lambda > 0$ and $\text{err}(\mathbf{x}, t) = 0$ ($x \geq 2R_A$), where c_R as well as λ depends only on R_A .

REMARK 3. The O terms in (3.23) or (3.24) depend on A , not only on R_A . If we write them in the form $\beta_A \times O(\cdot)$ with $\beta_A := m_{2R_A}(e_A)$, then the $O(\cdot)$ in the latter depend only on R_A . (One may realize this by observing that the dependence on A other than R_A comes in through (3.25) and (3.28) given below.) It is noted that $\sup_{\xi \in \partial U(2R_A)} e_A(\xi) < C\beta_A$ owing to Harnack's inequality, and that since $-\lg R + m_R(e_A)$ is independent of $R \geq R_A$, $\beta_A = \lg 2 + m_{R_A}(e_A)$. (See (6.13)).

Proof. The case $x \rightarrow \infty$ is dealt with in almost the same way as in the proof of Lemma 3.1 (except for $O(\cdot)$ term in (3.24)). Let $R = 2R_A$. We only notify that instead of the bound (3.9) we have

$$I_{[t/2, t]} \leq \frac{q(x, t; R)}{\lg x} \times c_R \beta_A e^{-x^2/3t} \quad \text{for } R < x < t. \quad (3.25)$$

We omit the proof since an upper bound of the left-hand side is computed in Appendix (Lemma 6.9), which is better than the bound above (according to Theorem 2.1).

In order to obtain the error terms $O(\cdot)$ of (3.23) and (3.24) we need to use the following refinement of (2.3): uniformly for $x > a$, as $t \rightarrow \infty$

$$\begin{aligned} q(x, t; a) &= \frac{4\pi \lg(x/a)}{(\lg t)^2} p_t^{(2)}(x) \left(1 + O\left(\frac{1}{\lg t}\right)\right) \quad \text{if } x < \sqrt{t}, \\ &= \frac{\pi}{\lg(t/x)} p_t^{(2)}(x) \left(1 + O\left(\frac{1}{\lg(t/x)}\right)\right) \quad \text{if } x/t \rightarrow 0, x \geq \sqrt{t}, \end{aligned} \quad (3.26)$$

which is a part of Corollary 4 of [25]. This allows us to replace $o(1)$ by $O(1/\lg(t/x))$ in (3.8) if $s/t \rightarrow 0$, so that taking $M = \sqrt{t}$ in the proof of Lemma 3.1 and employing (3.25) we deduce

$$q_A(\mathbf{x}, t) = q(x, t; R) \left(1 + O\left(\frac{1}{\lg(t/x)} \vee \frac{1}{\lg x}\right)\right) \quad \text{for } 2 \vee R < x < t/2, \quad (3.27)$$

which implies the second half of (3.23). Since $e_A(\mathbf{x}) = \lg x + O(1)$ (see (6.9)), it also shows (3.24) for $t^{1/4} \leq x < \sqrt{t}$ as well.

Finally consider the case $x < t^{1/4}$. We follow the proof of Theorem 3.1 in which we take $r = t^{1/3}$ and $T = t^{3/4}$. Observe that in the integral restricted to $[0, T]$ on the right-hand side of (3.13), $q_A(\xi, t - s)$ ($\xi \in \partial U(r)$) may be replaced by

$$\frac{4\pi \lg r}{(\lg t)^2} p_t^{(2)}(x) \left(1 + O\left(\frac{1}{\lg t}\right)\right)$$

in view of (3.26) and (3.27); also since $P_{\mathbf{x}}[\sigma_{\partial U(r)} \geq T] = O(e^{-\lambda T/r^2})$ for some $\lambda > 0$,

$$P_{\mathbf{x}}[\sigma_{\partial U(r)} < T \wedge \sigma_A] = P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] + O(e^{-\lambda t^{1/20}}).$$

Then, applying Lemma 3.3 below, we find that the integral over $[0, T]$ yields the formula on the right-hand side of (3.24) with $\text{err}(\mathbf{x}, t)$ removed. The integral over $[T, 3t/4]$ is $O(e^{-\lambda T/r^2})$ and that over $[3t/4, t]$ forms the error term $\text{err}(\mathbf{x}, t)$. Proof of Theorem 3.2 is complete. $\square$

Lemma 3.3. *For $r \geq 2R_A$ and for $\mathbf{x} \in \Omega_A \cap U(r)$,*

$$e_A(\mathbf{x}) \left(1 - \frac{3m_{R_A}(e_A)}{\lg(r/R_A)} \right) \leq P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \lg \left(\frac{r}{R_A} \right) \leq e_A(\mathbf{x}). \quad (3.28)$$

Here we write $m_R(e_A) = \int_{\partial U(R)} e_A(\xi) m_R(d\xi)$.

The proof as well as some variants of this lemma is given in Section 6.2.

3.1.3. ASYMPTOTIC FORM OF H_A AS $x/t \rightarrow 0$. Remember that $H_A(\mathbf{x}, t; d\xi)$ is defined in (1.2) for a bounded Borel set A .

Theorem 3.3. *Let $d \geq 3$. Uniformly for Borel subsets E of ∂A , as $t \rightarrow \infty$ and $x/t \rightarrow 0$*

$$H_A(\mathbf{x}, t; E) = q_A(\mathbf{x}, t)(H_A^\infty(E) + o(1)) + \text{err}(\mathbf{x}, t) \quad (3.29)$$

with $|\text{err}(\mathbf{x}, t)| \leq c_R t^{-d/2} P_{\mathbf{x}}[\sigma_{A \cup \partial U(2R_A)} > t/\lg t]$ (the same bound as in Theorem 3.1).

Proof. Let $R = 2R_A$ as before. We proceed as in the proof of Lemma 3.1. Suppose $x \rightarrow \infty$ and $x/t \rightarrow 0$. As before we write

$$H_A(\mathbf{x}, t; E) = \int_0^t ds \int_{\partial U(R)} H_{U(R)}(\mathbf{x}, t-s; d\xi) H_A(\xi, s; E) \quad (3.30)$$

and restrict the outer integral of (3.30) to $[0, M]$ with M to be made indefinitely large under $(1 \vee x^2)M/t^2 \rightarrow 0$, $M \leq \sqrt{t}$. For a probability measure m , write $P_m = \int m(d\xi) P_\xi$. Let $d \geq 3$ and observe $\text{Cap}(A) = \Lambda_\nu(0) R^{2\nu} P_{m_R}[\sigma_A < \infty]$ (see (3.2)) and substitute it into the formula of Theorem 3.1. Then in view of Theorem 2.1 we find

$$q_A(\mathbf{x}, t) = P_{\mathbf{x}}[\sigma_A = \infty] P_{m_R}[\sigma_A < \infty] q(x, t; R)(1 + o(1)), \quad (3.31)$$

in which the factor $P_{\mathbf{x}}[\sigma_A = \infty]$, tending to unity, can be dropped from the right-hand side. Now we apply Theorem 2.4 and (3.31) in turn. Noting $q(x, t-s; R) = q(x, t; R)(1 + o(1))$ for $s < M$, the integral in (3.30) restricted on $[0, M]$ then may be written as

$$\begin{aligned} & \int_0^M q(x, t-s; R) ds \int_{\partial U(R)} H_A(\xi, s; E) m_R(d\xi) (1 + O(x/t)) \\ &= q(x, t; R) P_{m_R}[B_{\sigma(A)} \in E, \sigma_A < M] (1 + o(1)) \\ &= q_A(\mathbf{x}, t) (P_{m_R}[B_{\sigma(A)} \in E \mid \sigma_A < \infty] + o(1)), \end{aligned} \quad (3.32)$$

of which the last expression may be further rewritten as the right-hand side of (3.29) since $P_{m_R}[B(\sigma_A) \in E \mid \sigma_A < \infty] = H_A^\infty(E)$. The remaining integral being dominated by one with $E = \partial A$, the required bounds of it obviously follows from those obtained in the proofs of Theorems 3.1.

The case when x remains in a bounded set can be dealt with in a similar way by following the corresponding part of the proof of Theorem 3.1. The details are omitted. $\square$

REMARK 4. Under the constraint $x < \sqrt{2t \lg t}$ the $o(1)$ term in (3.29) can be replaced by $O(t^{-\eta})$ with some constant $\eta > 0$, verification being the same as that of Remark 2 (c).

Theorem 3.4. *Let $d = 2$. Uniformly for Borel subsets E of ∂A , as $t \rightarrow \infty$ and $x/t \rightarrow 0$*

$$H_A(\mathbf{x}, t; E) = q_A(\mathbf{x}, t) \left(H_A^\infty(E) + O\left(\frac{1}{\lg[t/(t^{1/2} \vee x)]}\right) \right) + \text{err}(\mathbf{x}, t) \quad (3.33)$$

with $|\text{err}(\mathbf{x}, t)| \leq c_R t^{-1} (\lg t)^{-2} P_{\mathbf{x}}[\sigma_{A \cup \partial U(2R_A)} > t/2]$ (the same bound as in Theorem 3.2).

Proof. The proof is substantially the same as that of Theorem 3.3 apart from the $O(\cdot)$ term in (3.33). We consider only the case $R < x < t^{1/2}$ ($R = 2R_A$), the other case being similar. For $t^{1/4} < x < t^{1/2}$, according to Theorem 2.4 and (3.26) we have $q_A(\mathbf{x}, t) = q(x, t; R)(1 + O(1/\lg t))$ (in place of (3.31)) and observe that $O(x/t)$ and $o(1)$ in (3.32) can be replaced by $O((\lg t)^2 x/t)$ and $O(1/\lg t)$, respectively, which suffices for the required error estimate. For $x \leq t^{1/4}$, the argument given in the proof of Theorem 3.2 leads to the same error estimate. $\square$

In view of Theorem 2.4 the same argument as in the proof above shows

Lemma 3.4. *For $M > 0$ there exists a constant $c_M > 0$ such that if $R_A x/t < M$, $x > 2R_A$, and $t > R_A^2$, then*

$$H_A(\mathbf{x}, t; d\xi) \leq c_M q_A(\mathbf{x}, t) H_A^\infty(d\xi).$$

3.2 Case $x/t \rightarrow \infty$

In this subsection we suppose K to be a compact set. Let $\mathbf{e}$ be a unit vector and $\text{pr}_{\mathbf{e}} A$ the orthogonal projection of a set A on $\Delta_{\mathbf{e}}$, the hyper-plane perpendicular to $\mathbf{e}$ passing through the origin. We often write $\text{pr}_{\mathbf{x}}$ for $\text{pr}_{\mathbf{e}}$ if $\mathbf{e} = \mathbf{x}/x$. We bring in the subset $\langle K \rangle_{\mathbf{e}}$ of ∂K given by

$$\langle K \rangle_{\mathbf{e}} = \{\xi \in \partial K : \xi + t\mathbf{e} \notin K \text{ for } t > 0\}, \quad (3.34)$$

the mappings $h = h_{\mathbf{e}, K} : \Delta_{\mathbf{e}} \mapsto \mathbf{R}$ by the relations

$$\mathbf{z} + h(\mathbf{z})\mathbf{e} \in \langle K \rangle_{\mathbf{e}}, \quad \mathbf{z} \in \text{pr}_{\mathbf{e}} K,$$

and $h(\mathbf{z}) = -\infty$ for $\mathbf{z} \notin \text{pr}_{\mathbf{e}} K$, and the measure $m_{K, \mathbf{e}}$ on ∂K by

$$m_{K, \mathbf{e}}(E) = \text{vol}_{d-1}(\text{pr}_{\mathbf{e}}(\langle K \rangle_{\mathbf{e}} \cap E)) \quad (E \subset \partial K). \quad (3.35)$$

Here we regard $\text{pr}_{\mathbf{e}} K (= \text{pr}_{\mathbf{e}} \langle K \rangle_{\mathbf{e}})$ as a subset of $(d-1)$ -dimensional Euclidean space. Denote by $\partial_{d-1}(\text{pr}_{\mathbf{e}} K)$ the boundary of it as such. Note that $m_{K, \mathbf{e}}$ is supported by $\langle K \rangle_{\mathbf{e}}$. We need to further bring in the set of discontinuity points of h given by

$$\text{dis-ct}_{\mathbf{e}}(K) = \partial_{d-1}(\text{pr}_{\mathbf{e}} K) \cup \{\mathbf{z} \in \text{int}(\text{pr}_{\mathbf{e}} K) : h \text{ is discontinuous at } \mathbf{z}\}. \quad (3.36)$$

Theorem 3.5. *Let $\mathbf{e} \in \partial U(1)$ and suppose that $\text{dis-ct}_{\mathbf{e}}(K)$ is of zero Jordan measure, namely*

$$\text{vol}_{d-1}(\overline{\text{dis-ct}_{\mathbf{e}}(K)}) = 0. \quad (3.37)$$

Then, as $t \rightarrow \infty$ with $v := x/t \rightarrow \infty$

$$H_K(x\mathbf{e}, t; d\xi) \sim v p_t^{(d)}(x) e^{v\mathbf{e} \cdot \xi} m_{K, \mathbf{e}}(d\xi). \quad (3.38)$$

By (3.38) we mean that the Borel measure

$$\mu_{t,\mathbf{x}}^K(d\xi) := \frac{e^{-v\mathbf{e}\cdot\xi} H_K(x\mathbf{e}, t; d\xi)}{vp_t^{(d)}(x)} \quad (3.39)$$

($d\xi \subset \partial K$) weakly converges to $m_{K,\mathbf{e}}(d\xi)$, namely for any continuous φ on ∂K , the integral $\int \varphi(\xi) \mu_{t,\mathbf{x}}^K(d\xi)$ approaches $\int \varphi(\xi) m_{K,\mathbf{e}}(d\xi)$ (as $x/t \rightarrow \infty$). Note that $\xi \cdot \mathbf{e} \leq 0$, if $A \subset \{\mathbf{y} : \mathbf{e} \cdot \mathbf{y} \leq 0\}$. Although there is no direct way to derive from (3.38) any exact asymptotic form of $q_A(\mathbf{x}, t)$ and if the total mass $m_{K,\mathbf{e}}(\partial K)$ vanishes, (3.38) provides only a crude upper bound of $H_K(x\mathbf{e}; t, d\xi)$, the formula is useful for the study of Wiener sausage (see the end of Section 5) and of the hitting site distribution for the Brownian motion with large drift.

The formula (3.38) may be formally inferred by looking at the space-time distribution of the first arrival of (B_t) on the plane passing through ξ and perpendicular to $\mathbf{e}$. Indeed, if $B_0 = \mathbf{x}$, its density is given by

$$\frac{x - \xi \cdot \mathbf{e}}{t} p_t^{(1)}(x - \xi \cdot \mathbf{e}) p_t^{(d-1)}(|\text{pr}_{\mathbf{e}} \xi|), \quad (3.40)$$

which is asymptotic to the right-hand side of (3.38) divided by $m_{K,\mathbf{e}}(d\xi)$. The actual proof is postponed to the last part of Section 4.2, since we need to use a result given there.

REMARK 5. (a) There is a large class of K such that $\text{vol}_{d-1}(\text{pr}_{\mathbf{e}} K) = 0$ for all $\mathbf{e}$ and K is non-polar. This condition obtains if K has Hausdorff dimension larger than $d - 2$ and zero Hausdorff measure of $(d - 1)$ -dimension ([7] Theorem 6.4 ($d \geq 3$), [1] Exercise 27 of p.373 ($d = 2$)).

(b) It is not clear whether the condition (3.37) need to be required for (3.38). If $\langle K \rangle_{\mathbf{e}}$ is on a hyper-plane perpendicular to $\mathbf{e}$, it may be removed. For the upper bound the condition (3.37) is not needed. When a ball is taken for K an exact form of the density $H_{U(a)}(x\mathbf{e}, t; d\xi)/m_a(d\xi)$ is computed in [27], according to which it coincides with that in (3.38) except for a certain singularity appearing along the circumference of the hemisphere $\langle U(a) \rangle_{\mathbf{e}}$.

(c) For the question of (b) the presupposition that the set K is closed would be essential. As an example let G be an open set of the unit interval $[0, 1]$ with Lebesgue measure less than $1/2$ and containing all the rational number in $[0, 1]$ and F the complement of G in $[0, 1]$, and consider the set $A \subset \mathbf{R}^2$ defined by $A = [0, 1] \times G + \{0\} \times F$ (disjoint union). Then with $\mathbf{e} = (1, 0)$ we have $\langle A \rangle_{\mathbf{e}} = \{1\} \times G + \{0\} \times F$ but $H_A(x\mathbf{e}; t, d\xi)/q_A(\mathbf{x}, t)$ converges to the Lebesgue measure on $\{1\} \times [0, 1]$, so that the limit measure is not supported by $\langle A \rangle_{\mathbf{e}}$. By the way the compact set $K = [0, 1] \times F$ provides an example that does not satisfy (3.37) but $\text{vol}_{d-1}(\text{pr}_{\mathbf{e}} K) > 0$; indeed $\{0\} \times F = \langle K \rangle_{\mathbf{e}} = \partial \langle K \rangle_{\mathbf{e}}$ for the same $\mathbf{e}$. For this K the formula (3.38) holds (see Corollary 4.3).

Here we state and prove a simple lemma, which we shall apply in the proof of Proposition 4.1 given in the section 4.2.3.

Lemma 3.5. *There exists a constant $C > 1$ such that if $R_A x > t > 2R_A^2$, $E \subset \partial A$ and $H_A^\infty(E) > 0$,*

$$C^{-1} q(x, t - 1, R) P_{m_R}[B(\sigma_A) \in E, \sigma_A \leq 1] \leq H_A(\mathbf{x}, t; E) \leq C \beta_A q(x, t; R) H_A^\infty(E),$$

where $R = 2R_A$ and β_A equals 1 or $\text{Cap}(A)$ according as $d = 2$ or $d \geq 3$.

Proof. For $\xi \in \partial U(R)$, the ratio $\int_0^1 H_A(\xi, s; E) ds / P_{m_R}[B(\sigma_A) \in E, \sigma_A \leq 1]$ is bounded away from zero and infinity. Also it holds that $P_{m_R}[B(\sigma_A) \in E, \sigma_A \leq 1] \leq \beta_A H_A^\infty(E)$ and $q(x, t - 1; R) \leq 2q(x, t; R)$ (for t large enough). Using these inequalities we deduce the inequalities of the lemma from the Huygens formula (3.1). $\square$.

One can construct a compact set A such that the infimum of $P_{m_R}[B(\sigma_A) \in E, \sigma_A \leq 1] / \beta_A H_A^\infty(E)$ over Borel sets E with $H_A^\infty(E) > 0$ vanishes, so that in the lower bound in Lemma 3.5 $P_{m_R}[B(\sigma_A) \in E, \sigma_A \leq 1]$ cannot be replaced by $\beta_A H_A^\infty(E)$.

4 Wiener Sausage

Given a bounded Borel set A , let $S_A(t)$ be a Wiener sausage of length t swept by A attached to a Brownian motion B_t :

$$S_A(t) = \{\mathbf{z} \in \mathbf{R}^d : \mathbf{z} - B_s \in A \text{ for some } s \in [0, t]\}.$$

The d -dimensional volume of A is denoted by $\text{vol}_d(A)$. We sometimes write $\text{area}(A)$ for $\text{vol}_2(A)$; also we usually suppress (d) from $p_t^{(d)}$ and the subscript $\mathbf{0}$ from $P_{\mathbf{0}}$ and $E_{\mathbf{0}}$. Throughout this section we suppose that A is non-polar.

4.1 Case $\mathbf{x}/t \rightarrow 0$

Theorem 4.1. *If $d \geq 3$, uniformly for $\mathbf{x}$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$,*

$$E_{\mathbf{0}}[\text{vol}_d(S_A(t)) \mid B_t = \mathbf{x}] \sim \text{Cap}(A)t. \quad (4.1)$$

Theorem 4.2. *If $d = 2$, uniformly for $\mathbf{x}$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$,*

$$E_{\mathbf{0}}[\text{area}(S_A(t)) \mid B_t = \mathbf{x}] = \begin{cases} \frac{2\pi t}{\lg t}(1 + o(1)) & \text{if } x \leq \sqrt{t}, \\ \frac{\pi t}{\lg(t/x)}(1 + o(1)) & \text{if } x > \sqrt{t}. \end{cases} \quad (4.2)$$

REMARK 6. (a) In the case when B_t is pinned at $\mathbf{x} = \mathbf{0}$ an asymptotic expansion of $E_{\mathbf{0}}[\text{vol}_d(S_A(t)) \mid B_t = \mathbf{0}]$ is obtained in [15] ($d \geq 3$) and [16] ($d = 2$) (see also [3]).

(b) In [24], it is shown that if $d = 2$, for each $M > 1$, uniformly for $|\mathbf{x}| \leq M\sqrt{t}$,

$$E_{\mathbf{0}}[\text{area}(S_{U(a)}(t)) \mid B_t = \mathbf{x}] = 2\pi t N(\kappa t/a^2) + \frac{\pi x^2}{(\lg t)^2} \left[\lg \frac{t}{x^2 \vee 1} + O(1) \right] + O(1)$$

as $t \rightarrow \infty$, where $\kappa = 2e^{-2\gamma}$ and $N(\lambda) = \int_0^\infty e^{-\lambda u} [(\lg u)^2 + \pi^2]^{-1} u^{-1} du$, $N(\alpha t) = \frac{1}{\lg t} - \frac{\gamma + \lg \alpha}{(\lg t)^2} + \dots$ (asymptotic expansion in powers of $1/\lg t$).

(c) The corresponding problem for pinned random walk on $\mathbb{Z}^2$ is studied in [22] and [28].

Lemma 4.1. *For $0 < t_1 \leq t$, $\mathbf{x} \in \mathbf{R}^d$ and a Borel set $W \subset \mathbf{R}^d$ such that $A \cap W = \emptyset$,*

$$\begin{aligned} & E_{\mathbf{0}}[\text{vol}_d(S_A(t_1) \cap W) \mid B_t = \mathbf{x}] \\ &= \frac{1}{p_t(x)} \int_{\mathbf{z} \in W} |d\mathbf{z}| \int_0^{t_1} ds \int_{\xi \in \partial A} p_{t-s}(\mathbf{z} - \mathbf{x} - \xi) H_A(\mathbf{z}, s : d\xi), \end{aligned} \quad (4.3)$$

where $|\cdot|$ designates the Lebesgue measure on $\mathbf{R}^d$. In particular, putting

$$F_A(t, \mathbf{x}) = \int_{\mathbf{z} \notin A} |d\mathbf{z}| \int_0^t ds \int_{\xi \in \partial A} p_{t-s}(\mathbf{z} - \mathbf{x} - \xi) H_A(\mathbf{z}, s : d\xi)$$

we have

$$E_0[\text{vol}_d(S_A(t)) | B_t = \mathbf{x}] p_t(\mathbf{x}) = F_A(t, \mathbf{x}) + \text{vol}_d(A) p_t(\mathbf{x}). \quad (4.4)$$

Proof. This proof is standard but we need it later (see the second half of proof of Lemma 4.4). Let the Brownian motion B_t be started at the origin. Let $-A + \mathbf{z}$ denote the set $\{-\xi + \mathbf{z} : \xi \in A\}$. Then noting that for any $\varepsilon > 0$

$$S_A(t_1) \cap W \subset \{\mathbf{z} \in W : \sigma_{-A+\mathbf{z}} \leq t_1\} \subset S_A(t_1 + \varepsilon) \cap W, \quad (4.5)$$

we see that

$$\begin{aligned} & E[\text{vol}_d(S_A(t_1) \cap W) | B_t = \mathbf{x}] p_t(x) \\ &= \int_W P[\sigma_{-A+\mathbf{z}} \leq t_1 | B_t = \mathbf{x}] p_t(x) |d\mathbf{z}| \\ &= \int_W |d\mathbf{z}| \int_{[0, t_1] \times \partial(-A+\mathbf{z})} p_{t-s}(\mathbf{x} - \xi) P[\sigma_{-A+\mathbf{z}} \in ds, B(\sigma_{-A+\mathbf{z}}) \in d\xi] \\ &= \int_W |d\mathbf{z}| \int_0^{t_1} ds \int_{\xi \in -\partial A} p_{t-s}(\mathbf{x} - \mathbf{z} - \xi) H_{-A}(-\mathbf{z}, s; d\xi), \end{aligned}$$

which, owing to the equality $\int_{-\partial A} \varphi(\xi) H_{-A}(-\mathbf{z}, s; d\xi) = \int_{\partial A} \varphi(-\xi) H_A(\mathbf{z}, s; d\xi)$, is the same as the formula to be shown. $\square$

Put

$$F_A^*(t, \mathbf{x}) = \int_{\mathbf{z} \notin A} |d\mathbf{z}| \int_0^t p_{t-s}(\mathbf{z} - \mathbf{x}) ds \int_{\xi \in \partial A} e^{-\mathbf{x} \cdot \xi / t - |\xi|^2 / 2t} H_A(\mathbf{z}, s : d\xi).$$

By the scaling property of Brownian motion we have

$$F_A(t, \mathbf{x}) = F_{A/r}(t/r^2, \mathbf{x}/r), \quad (4.6)$$

and similarly for F_A^* .

Lemma 4.2. *Let M be a positive constant. If $d \geq 3$, for $R_A x/t < M$ and $t > R_A^2$,*

$$|F_A(t, \mathbf{x}) - F_A^*(t, \mathbf{x})| \leq c_M p_t(x) \left[R_A^{2\nu} + \text{Cap}(A) \sqrt{t}/R_A \right],$$

where c_M is a constant depending only on M and d . If $d = 2$, the same bound with $\text{Cap}(A)$ replaced by unity holds true.

Proof. The proof is similar to that of Lemma 4.3 of [24]. Let $a > 4$ be a constant that is suitably chosen later depending on M (see the part (c) below). Putting $T = s(t-s)/t$, we bring in the following sub-regions of $D := [0, t] \times \mathbf{R}^d$

$$\begin{aligned} D^a &= \{(s, \mathbf{z}) : t-a < s \leq t, \mathbf{z} \in \Omega_A\}, \\ D_a &= \{(s, \mathbf{z}) : 0 < s \leq a, \mathbf{z} \in \Omega_A\}, \\ D_{>} &= \{(s, \mathbf{z}) : a < s \leq t-a, |\mathbf{z} - (s/t)\mathbf{x}| \geq \sqrt{8T \lg T}, \mathbf{z} \in \Omega_A\}, \\ D_{<} &= \{(s, \mathbf{z}) : a < s \leq t-a, |\mathbf{z} - (s/t)\mathbf{x}| < \sqrt{8T \lg T}, \mathbf{z} \in \Omega_A\}, \end{aligned}$$

and restrict to them the integrals that define F_A or F_A^* . Denote the corresponding integrals for F_A by $J\{D^a\}$, $J\{D_a\}$, etc. and those for F_A^* by $J^*\{D^a\}$, $J^*\{D_a\}$, etc.

Of the first three regions above we evaluate the corresponding integrals J and J^* separately, and verify that they are all bounded from above by

$$c_M R_A^{2\nu} p_t(x)$$

in the paragraphs (a), (b) and (c) below. The actual computations are given only for F_A , those for F_A^* being similar and much simpler. From Proposition 6.3 we have

$$q_A(\mathbf{z}, t) \leq c_M \text{Cap}(A) p_t(|\mathbf{z}|) e^{R_A |\mathbf{z}|/t} \quad \text{for } t > R_A^2, \mathbf{z} \in \Omega_A, \quad (4.7)$$

For simplicity let $R_A = 1$ in what follows, which gives no loss of generality because of the scaling property (4.6), and suppose $t > 2a$ and $x < Mt$. Denote by c_M, c'_M etc. unimportant positive constants that depends only on M and d .

(a) D^a : Use the inequality

$$H_A(\mathbf{z}, s; d\xi) \leq \int_{\Omega_A} p_1(\mathbf{y} - \mathbf{z}) H_A(\mathbf{y}, s-1; d\xi) |d\mathbf{y}| \quad (s > 1)$$

and perform integration by $\mathbf{z}$ first. Then, after changing variable of integration, we have

$$J\{D^a\} \leq \int_{\Omega_A} |d\mathbf{y}| \int_0^a ds \int_{\partial A} p_{s+1}(\mathbf{y} - \mathbf{x} - \xi) H_A(\mathbf{y}, t-s-1; d\xi).$$

Using the trivial bound $p_{s+1}(\cdot) \leq 1$ for $|\mathbf{y}| > 2x$ and the bound given in Lemma 3.4 of $H_A(\cdot, \cdot; d\xi)/H_A^\infty(d\xi)$ for $|\mathbf{y}| < 2x$ together with (4.7) we evaluate the repeated integral to be at most

$$\text{Cap}(A) \left[o(p_t(x)) + c_M \int_{\partial A} p_t(\mathbf{x} - \xi) H_A^\infty(d\xi) \right] \leq c'_M \text{Cap}(A) p_t(x)$$

as desired, where we have applied the assumption $x < Mt$ for the last inequality.

(b) D_a : Noting $\sup_{\xi \in \bar{A}} p_{t-s}(\mathbf{z} - \mathbf{x} - \xi) < \kappa_d p_{t+1}(\mathbf{z} - \mathbf{x})$ ($s < a$) and $\sigma_A \geq \sigma_{U(1)}$ we see

$$J\{D_a\} \leq \kappa_d \int_{\Omega_d} p_{t+1}(\mathbf{z} - \mathbf{x}) P_{\mathbf{z}}[\sigma_A < a] |d\mathbf{z}| \leq \kappa_d p_{t+a+2}(x) \leq c_M p_t(x),$$

where we have applied Theorem 2.2 for the second inequality.

For the rest of this proof we shall use the identity

$$p_s(\mathbf{z}) p_{t-s}(\mathbf{z} - \mathbf{x}) = p_t(\mathbf{x}) p_T\left(\mathbf{z} - \frac{s}{t} \mathbf{x}\right) \quad (4.8)$$

(the second factor on the right side is the probability density of the Brownian bridge at time s); remember

$$T = T(t, s) = \frac{s(t-s)}{t}.$$

(c) $D_{>}$: Using the inequalities $p_s \leq \kappa_d p_{s+1}$ and $p_s(x) e^{x/s} \leq \kappa_d p_{s+1}(x)$ ($s > 1$) as well as (4.7) we have

$$J\{D_{>}\} \leq c_M \text{Cap}(A) \int_a^{t-a} ds \int_{|\mathbf{z} - (s/t)\mathbf{x}| > \sqrt{8T \lg T}} p_{s+1}(\mathbf{z}) p_{t-s+1}(\mathbf{z} - \mathbf{x}) |d\mathbf{z}|.$$

On writing $t' = t+2$, $s' = s+1$ and $T' = s'(t' - s')/t'$ the integral on the right-hand side divided by $p_{t+2}(x)$ equals $\int_a^{t-a} ds \int_{|\mathbf{z} - (s/t)\mathbf{x}| > \sqrt{8T \lg T}} p_{T'}(\mathbf{z} - \frac{s'}{t'}\mathbf{x}) |d\mathbf{z}|$. Noting $|s'/t' - s/t|x \leq x/t < M$, choose $a = a_M$ so that $\sqrt{8T \lg T} - M \geq \sqrt{5T' \lg T'}$ for $a < s < t - a$. We can then conclude that

$$\frac{J\{D_{>}\}}{\text{Cap}(A)p_t(x)} \leq c_M \int_a^{t-a} ds \int_{\sqrt{5 \lg T'}} e^{-u^2/2} u^{d-1} du \leq c'_M \int_{a+1}^{t'-a} \frac{ds'}{[s'(t' - s')/t']^2} \leq c''_M a^{-1}$$

as required.

(d) $D_{<}$: Let $R_A = 1$ as above. Using (4.8) we observe

$$\begin{aligned} & \frac{p_s(\mathbf{z})}{p_t(\mathbf{x})} \left[p_{t-s}(\mathbf{z} - \mathbf{x} - \xi) - e^{-\mathbf{x} \cdot \xi / t - |\xi|^2 / 2t} p_{t-s}(\mathbf{z} - \mathbf{x}) \right] \\ &= \frac{p_t(\mathbf{x} + \xi)}{p_t(\mathbf{x})} \left[p_T\left(\mathbf{z} - \frac{s}{t}(\mathbf{x} + \xi)\right) - p_T\left(\mathbf{z} - \frac{s}{t}\mathbf{x}\right) \right]. \end{aligned} \quad (4.9)$$

For $(s, \mathbf{z}) \in D_{<}$ we have $|\mathbf{z} - \frac{s}{t}\mathbf{x}|/T \leq \sqrt{8(\log T)/T} < 3$ and, on using $|e^{\alpha-p} - 1| \leq e^{|\alpha|}(|\alpha| + p)$ ($p \geq 0$),

$$\begin{aligned} & \left| p_T\left(\mathbf{z} - \frac{s}{t}(\mathbf{x} + \xi)\right) - p_T\left(\mathbf{z} - \frac{s}{t}\mathbf{x}\right) \right| \\ &= p_T\left(\mathbf{z} - \frac{s}{t}\mathbf{x}\right) \left| \exp \left\{ \frac{s/t}{T} \left[\left(\mathbf{z} - \frac{s}{t}\mathbf{x}\right) \cdot \xi - \frac{s|\xi|^2}{2t} \right] \right\} - 1 \right| \\ &\leq e^3 p_T\left(\mathbf{z} - \frac{s}{t}\mathbf{x}\right) \frac{1}{t-s} \left[|\xi| \left| \mathbf{z} - \frac{s}{t}\mathbf{x} \right| + \frac{s|\xi|^2}{2t} \right]. \end{aligned} \quad (4.10)$$

Hence, if $x \leq Mt$,

$$\begin{aligned} & |J\{D_{<}\} - J^*\{D_{<}\}| \\ &\leq e^3 e^M \int_{D_{<}} \frac{p_t(\mathbf{x})}{p_s(\mathbf{z})} \cdot p_T\left(\mathbf{z} - \frac{s}{t}\mathbf{x}\right) \frac{1}{t-s} \left(\left| \mathbf{z} - \frac{s}{t}\mathbf{x} \right| + \frac{s}{2t} \right) q_A(\mathbf{z}, s) ds |d\mathbf{z}|. \end{aligned}$$

On $D_{<}$ we have $|\mathbf{z}| < (M+1)s$ and, noting $\int p_T(\mathbf{y})|y||d\mathbf{y}| \leq \kappa_d \sqrt{T}$, we apply (4.7) to deduce that the integral above is at most a constant multiple of

$$\text{Cap}(A)p_t(x) \int_a^{t-a} \frac{1}{(t-s)} \left[\sqrt{T} + \frac{s}{t} \right] ds \leq C' \text{Cap}(A)p_t(x) \sqrt{t}. \quad (4.11)$$

This completes the proof of Lemma 4.2. $\square$

Proof of Theorems 4.1 and 4.2. Owing to Lemma 4.2 as well as (4.4) it suffices to show that as $x/t \rightarrow 0$ and $t \rightarrow \infty$ the function $F_A^*(t, \mathbf{x})$, which may obviously be written as

$$\int_0^t ds \int_{\mathbf{z} \in \Omega_A} p_{t-s}(\mathbf{z} - \mathbf{x}) q_A(\mathbf{z}, s) |d\mathbf{z}| (1 + O(x/t)), \quad (4.12)$$

has the same asymptotic form as given on the right-hand side of (4.1) if $d \geq 3$ and that of (4.2) if $d = 2$. Put $D_t = \left\{ (s, \mathbf{z}) : 4 < s < t/2, |\mathbf{z} - (s/t)\mathbf{x}| < \sqrt{2s \lg s}, \mathbf{z} \notin A \right\}$.

Let $d \geq 3$. Considering D_t in place of $D_<$ we argue as in the proof of Lemma 4.2. Observe $\int_{|\mathbf{z}-(s/t)\mathbf{x}| \geq \sqrt{2s \lg s}} p_T(\mathbf{z} - \frac{s}{t}\mathbf{x}) |d\mathbf{z}| = O(1/s)$, $s > 1$ (valid even if s, t are replaced by $s+1, t+1$) and substitute the formula for q_A in Theorem 3.1 to see that the inner integral restricted to D_t equals

$$\begin{aligned} & \text{Cap}(A) p_t(\mathbf{x}) \int_{|\mathbf{z}-(s/t)\mathbf{x}| < \sqrt{2s \lg s}} P_{\mathbf{z}}[\sigma_A = \infty] p_T\left(\mathbf{z} - \frac{s}{t}\mathbf{x}\right) |d\mathbf{z}| (1 + o(1)) \\ &= \text{Cap}(A) p_t(\mathbf{x}) (1 + o(1)). \end{aligned}$$

Hence the integral on the lower half interval $0 < s < t/2$ gives half the asserted leading term in Theorem 4.1. The other half is dealt with in an analogous way and the details are omitted.

The case $x < M\sqrt{t}$ of $d = 2$ follows from the result for $U(a)$ given in Remark 6 (b) on noting that q_A has the same asymptotic form as $q_{U(1)}$ (for large space variable) in view of Theorems 2.1 and 3.2 and that the repeated integral (4.12) restricted on the outside of D_t is negligible. As for the other case of $d = 2$, of which the proof is somewhat involved if argued as above, we apply Proposition 4.1 of the next subsection, which immediately implies the asserted result in view of Theorem 3.2. $\square$

4.2 Case $x/\sqrt{t} \rightarrow \infty$

Here we consider the case $x/\sqrt{t} \rightarrow \infty$, mostly the case $x/t \rightarrow \infty$. The main results are presented in the first part of this subsection and proofs of them will be given in the second through fourth parts. Throughout this subsection we fix a unit vector $\mathbf{e} \in \partial U(1)$.

4.2.1. STATEMENTS OF RESULTS. Combined with Theorems 3.1 and 3.2 the next proposition covers the case $x/\sqrt{t} \rightarrow \infty$ of Theorems 4.2 as noted at the end of the preceding subsection. It also plays a substantial role in the proof of Theorem 3.5 and Theorem 4.3 below. A continues to denote a bounded Borel set of $\mathbf{R}^d$.

Proposition 4.1. *As $x^2/t \rightarrow \infty$ and $t \rightarrow \infty$*

$$\begin{aligned} E_{\mathbf{0}} [\text{vol}_d(S_A(t)) \mid B_t = \mathbf{x}] &\sim \frac{t}{p_t^{(d)}(x)} \cdot \frac{E_{\mathbf{x}}[e^{-B_{\sigma(A)} \cdot \mathbf{x}/t}; \sigma_A \in dt]}{dt} \\ &= \frac{t q_A^{(d)}(x, t)}{p_t^{(d)}(x)} \int_{\partial A} e^{-\xi \cdot \mathbf{x}/t} P_{\mathbf{x}}[B_{\sigma(A)} \in d\xi \mid \sigma_A = t]. \end{aligned}$$

Theorem 4.3. *Suppose that K is a compact set of $\mathbf{R}^d$ ($d \geq 2$), $\mathbf{e} \in \partial U(1)$ and the condition (3.37) is satisfied. Let $\mathbf{x} = x\mathbf{e}$. Then as $x/t \rightarrow \infty$ and $t \rightarrow \infty$,*

$$E_{\mathbf{0}} [\text{vol}_d(S_K(t)) \mid B_t = \mathbf{x}] \sim \text{vol}_{d-1}(\text{pr}_{\mathbf{e}} K) x. \quad (4.13)$$

The proofs of these two results and that of Theorem 3.5 are interrelated in a way. Our proof of Proposition 4.1 is quite involved: the difficulty occurs when x/t becomes indefinitely large, the proof being much easier otherwise. The upper bound for Theorem 4.3 is relatively easy and the result is useful for the proof of Proposition 4.1 in the case $x/t \rightarrow \infty$. For the lower bound of some cases of K of special shape is easy and it together with the upper bound

is used for the proof of Theorem 3.5, which combined with Proposition 4.1 immediately gives Theorem 4.3.

In the rest of this part we give a proof of (vi), which concerns the case when x/t approaches a positive constant v and need Proposition 4.1. We remind the reader that for each $a > 0$, the function

$$g_{av}(\theta) = \sum_{n=0}^{\infty} \frac{K_0(av)}{K_n(av)} H_n(\theta)$$

is a probability density on $\partial U(1)$ with respect to $m_1(d\xi)$. Here $\theta = \theta(\xi)$ is the colatitude of ξ relative to an vector $\mathbf{e} \in \partial U(1)$, which may be arbitrarily taken as the north pole (see (2.5)).

Proof of (vi) of Section 1.

The second formula of (vi) follows immediately from the first owing to Proposition 4.1. As for the first one we recall the Huygens decomposition (3.1) and substitute the formulae of Theorems 2.1 and 2.4 into it. Write $\tilde{v} = x/t$. Noting that $p_{t-s}(\mathbf{x} - \xi)$ is decreasing in s whenever $0 < t - s < d^{-1}|\mathbf{x} - \xi|^2$ and that for $\xi \in \partial U(R)$,

$$p_{t-\sqrt{t}}(x - \xi)/p_t(x) \leq (1 - \frac{1}{\sqrt{t}})^{-d/2} e^{\xi \cdot \mathbf{x}/t} e^{-|\mathbf{x} - \xi|^2/2t\sqrt{t}} \leq c_M e^{-\frac{1}{2}\tilde{v}^2\sqrt{t}}$$

for $\tilde{v} = x/t < M$, we then infer that the main contribution to the outer integral in (3.1) comes from $[0, \sqrt{t}]$, so that as $x/t \rightarrow v$

$$\begin{aligned} & H_A(\mathbf{x}, t; E)/p_t(x) \\ &= \frac{R_A^{2\nu} \Lambda_\nu(R_A \tilde{v})}{p_t(x)} \int_0^{\sqrt{t}} p_{t-s}(x) ds \int_{\partial U(1)} H_A(R_A \xi, s; E) g_{R_A v}(\theta) m_1(d\xi) + \varepsilon_{t,x} \end{aligned}$$

with $\varepsilon_{t,x} \leq c_M \beta_A e^{\tilde{v}^2\sqrt{t}} R_A^{2\nu} \Lambda(R_A v)$, where θ is the colatitude of ξ relative to $\mathbf{e} = \mathbf{x}/x$ and β_A equals 1 or $\text{Cap}(A)$ according as $d = 2$ or $d \geq 3$ as in Lemma 3.5. Since $p_{t-s}(x) \sim p_t(x) e^{-\tilde{v}^2 s/2}$, on putting

$$\begin{aligned} & \lambda_A(\mathbf{v}; E) \\ &= R_A^{2\nu} \int_0^\infty e^{-v^2 s/2} ds \int_{\partial U(1)} H_A(R_A \xi, s; E) g_{R_A v}(\theta) m_1(d\xi) \end{aligned} \tag{4.14}$$

for $v = |\mathbf{v}| > 0$, this yields the formula $H_A(\mathbf{x}, t; E) \sim p_t(x) \Lambda_\nu(R_A \tilde{v}) \lambda_A(\mathbf{x}/t; E)$ as in (vi). Since $\lambda_A(\mathbf{v}; \partial A) \sim q_A(\mathbf{x}, t)/p_t(x) \Lambda_\nu(R_A v)$, in view of Propositions 6.3 and 6.4

$$\lambda_A(\mathbf{v}; \partial A) \leq c_M R_A^{2\nu} \beta_A.$$

We may define $\lambda_A(\mathbf{0}; d\xi)$ by

$$\lambda_A(\mathbf{0}; d\xi) = \int_0^\infty ds \int_{\partial U(1)} H_A(R_A \xi, s; E) m_1(d\xi) = \begin{cases} H_A^\infty(d\xi) & (d = 2) \\ G^{(d)}(1) \text{Cap}(A) H_A^\infty(d\xi) & (d \geq 3) \end{cases}$$

so as to have $\lambda_A(\mathbf{v}; \cdot)$ continuous at $\mathbf{v} = \mathbf{0}$; one then observes that the two asymptotic formulae of (vi) are valid uniformly about $v = 0$ and in particular they recover the results in the regime $x/t \rightarrow 0$ given in (iii), (iv) and the second half of (v). The asserted uniformity of the convergence in total variation norm is easily checked.

The formula (1.9) is an immediate consequence of Theorem 2.4 and Proposition 4.1.

4.2.2. UPPER BOUND FOR THEOREM 4.3. Here we prove the upper bound, which is reduced to the next proposition; this entails that K may be any bounded Borel set for the upper bound. The proof of the lower bound is postponed to Section 4.2.4.

Lemma 4.3. *For $a > 0$ and $h > 0$ put $C = \{\mathbf{z} : 0 \leq \mathbf{z} \cdot \mathbf{e} \leq h \text{ and } |\text{pr}_{\mathbf{e}} \mathbf{z}| \leq a\}$ (a circular cylinder of height h , radius a and axis $\mathbf{e}$). Then, as $v := x/t \rightarrow \infty$, $t \rightarrow \infty$,*

$$(i) \quad E_{\mathbf{0}}[\text{vol}_d(S_C(t)) \mid B_t = x\mathbf{e}] = c_{d-1}a^{d-1}x + O\left(a^{d-2}t\sqrt{hv} + (a^{d-1} + h)\sqrt{t}\right)$$

uniformly in a and h , and

$$(ii) \quad E_{\mathbf{0}}[(\text{vol}_d(S_C(t)))^2 \mid B_t = x\mathbf{e}] = [c_{d-1}a^{d-1}x]^2(1 + o(1)),$$

where $o(1)$ depends only on a , h and d and c_n denotes the volume of n -dimensional unit ball.

Proof. We give a proof only for $d = 2$, the higher dimensional case being essentially the same. Let $\mathbf{e} = \mathbf{e}_0 = (1, 0)$ and write B'_t and B''_t for the first and the second component of B_t so that $B_t = (B'_t, B''_t)$. For the first assertion (i) the lower bound is obvious: the expectation in it is bounded below by $c_{d-1}a^{d-1}x$. To see the upper bound let $\varepsilon > 0$ and put $N = \lceil t/\varepsilon \rceil$ and $t_k = kt/N$, so that $\varepsilon(1 - \varepsilon/t) < t_{k+1} - t_k \leq \varepsilon$. Noting that the conditional law of (B_s) given $B_t = x\mathbf{e}_0$ equals that of $(B_s + v\mathbf{e}_0)$ given $B_t = \mathbf{0}$, we bring in the variables

$$\xi_k^+ = \sup_{t_k \leq s \leq t_{k+1}} (B'_s + vs), \quad \xi_k^- = \inf_{t_k \leq s \leq t_{k+1}} (B'_s + vs); \text{ and}$$

$$\eta_k = \sup_{t_k \leq s \leq t_{k+1}} |B''_s - B''_{t_k}|.$$

Write $\Delta_k S$ for $S_C(t_{k+1}) \setminus S_C(t_k)$ and $Y_k^\pm = B''_{t_k} \pm a$. We decompose $\Delta_k S$ into two parts $\Delta_k S \setminus (\mathbf{R} \times [Y_k^-, Y_k^+])$ and $\Delta_k S \cap (\mathbf{R} \times [Y_k^-, Y_k^+])$. For the first part we have

$$\Delta_k S \setminus (\mathbf{R} \times [Y_k^-, Y_k^+]) \subset \left([\xi_k^-, \xi_k^+ + h] \times ([Y_k^+, Y_k^+ + \eta_k] \cup [Y_k^- - \eta_k, Y_k^-]) \right)$$

of which the volume of the right hand side is at most

$$2(h + \xi_k^+ - \xi_k^-)\eta_k. \tag{4.15}$$

As for the second part note that $\xi_k^- \leq \xi_{k-1}^+$ and $|Y_k^+ - Y_{k-1}^+| \vee |Y_k^- - Y_{k-1}^-| \leq \eta_{k-1}$ and that $S_C(t_k)$ includes the rectangle

$$[\xi_{k-1}^-, \xi_{k-1}^+ + h] \times [Y_k^- + 2\eta_{k-1}, Y_k^+ - 2\eta_{k-1}]$$

($[s, t]$ is understood to be the empty set if $s > t$), and one then deduces that

$$\begin{aligned} & \Delta_k S \cap (\mathbf{R} \times [Y_k^-, Y_k^+]) \\ & \subset ([\xi_{k-1}^+ + h, \xi_k^+ + h] \times [Y_k^-, Y_k^+]) \cup ([\xi_{k-1}^- \wedge \xi_k^-, \xi_{k-1}^-] \times [Y_k^-, Y_k^+]) \\ & \quad \cup ([\xi_{k-1}^-, \xi_{k-1}^+ + h] \times ([Y_k^+ - 2\eta_{k-1}, Y_k^+] \cup [Y_k^-, Y_k^- + 2\eta_{k-1}])) \end{aligned}$$

($k = 1, \dots, N-1$) and $\Delta_0 S \cap (\mathbf{R} \times [Y_0^-, Y_0^+]) \subset [\xi_0^-, \xi_0^+] \times [-a, a]$. Combined with the bound (4.15) this shows

$$\begin{aligned} \text{area}(S_C(t)) &\leq \sum_{k=0}^{N-1} 2a(\xi_k^+ - \xi_{k-1}^+) + 6 \sum_{k=0}^{N-1} (h + \xi_k^+ - \xi_k^-) \eta_k \\ &\quad + 2a \sum_{k=0}^{N-1} (\xi_k^- - \xi_{k+1}^-) \vee 0, \end{aligned} \quad (4.16)$$

where ξ_{-1}^+ is understood to be ξ_0^- . The expectations under the conditional measure given $B_t = \mathbf{0}$ of the variables $\xi_k^+ - \xi_k^-$ and η_k are at most constant multiples of $v\varepsilon + \sqrt{\varepsilon} + \varepsilon/\sqrt{t}$ and of $\sqrt{\varepsilon} + \varepsilon/\sqrt{t}$, respectively, for every k : indeed,

$$E[\eta_k \mid B_t = \mathbf{0}] \leq E\left[\sup_{t_k \leq s \leq t_{k+1}} |B_s'' - B_{t_k}''| + \varepsilon t^{-1} |B_t''|\right] < C\sqrt{\varepsilon} + C'\varepsilon/\sqrt{t}$$

and similarly for $\xi_k^+ - \xi_k^-$. Also $E[\xi_{N-1}^+] \leq x + E[\max_{0 \leq s \leq t} B_s'] \leq x + C\sqrt{t}$. Now we take $\varepsilon = h/v$. The processes (ξ_k^+, ξ_k^-) and (η_k) being independent under the law of the Brownian bridge, we see

$$E\left[\text{area}(S_C(t)) \mid B_t = x\mathbf{e}_0\right] \leq 2ax + O(a\sqrt{t}) + O(t\sqrt{hv}) + O(h\sqrt{t}).$$

This proves (i). The assertion (ii) follows from (4.16) as is easily checked. $\square$

From the proof of Lemma 4.3 its first assertion (i) may be slightly generalized so that for each $\delta \in (0, 1]$,

$$E_0[\text{vol}_d(S_C(\delta t)) \mid B_t = x\mathbf{e}] = c_{d-1}a^{d-1}\delta x + o(\delta x).$$

We shall need the upper bound which may be stated as follows.

Corollary 4.1. *Let K be a compact set of $\mathbf{R}^d$. Then for each $\delta \in (0, 1]$, as $x/t \rightarrow \infty$ and $t \rightarrow \infty$, $E_0[\text{vol}_d(S_K(\delta t)) \mid B_t = x\mathbf{e}] \leq \text{vol}_{d-1}(\text{pr}_{\mathbf{e}}K)\delta x + o(\delta x)$.*

Proof. The proof is given only in the case $\delta = 1$. Given $\varepsilon > 0$, we can find a finite number of balls $b_n \subset \Delta_{\mathbf{e}}, n = 1, \dots, N$ such that $\text{pr}_{\mathbf{e}}K \subset \cup b_n$ and $\sum \text{vol}_{d-1}(b_n) < \text{vol}_{d-1}(\text{pr}_{\mathbf{e}}K) + \varepsilon$. According to Lemma 4.3 we have that $E_0[\text{vol}_d(S_K(t)) \mid B_t = x\mathbf{e}]/x \leq \sum_{n=1}^N \left(\text{vol}_{d-1}(b_n) + O(1/\sqrt{v}) \right)$, where $O(1/\sqrt{v})$ is independent of b_n , showing the asserted inequality of the lemma. $\square$

Corollary 4.2. *Let K be a compact set of $\mathbf{R}^d$ such that K lies on a plane perpendicular to $\mathbf{e}$. Then as $x/t \rightarrow \infty$ and $t \rightarrow \infty$, $E_0[\text{vol}_d(S_K(t)) \mid B_t = x\mathbf{e}] \sim \text{vol}_{d-1}(K)x$.*

Proof. The lower bound holds path-wise due to the hypothesis on K , whereas the upper bound follows from Corollary 4.1. $\square$

4.2.3. PROOF OF PROPOSITION 4.1. Let A be a bounded Borel set of $\mathbf{R}^d$ ($d \geq 2$). In terms of H_A Proposition 4.1 may be stated as follows:

$$E_0[\text{vol}_d(S_A(t)) \mid B_t = \mathbf{x}] = \int_{\xi \in \partial A} e^{-\mathbf{x} \cdot \xi/t} H_A(\mathbf{x}, t; d\xi) \frac{t}{p_t(x)} (1 + o(1)) \quad (4.17)$$

as $x/\sqrt{t} \rightarrow \infty$ and $t \rightarrow \infty$. The proof is performed by showing Lemmas 4.4 and 4.5 given below. Put for $4 < t_1 \leq t$

$$J(t_1) = \frac{1}{p_t(x)} \int_{\mathbf{z} \in \Omega_A} |d\mathbf{z}| \int_4^{t_1-4} p_{t-s}(\mathbf{x} - \mathbf{z}) ds \int_{\partial A} e^{-\mathbf{x} \cdot \xi/t} H_A(\mathbf{z}, s; d\xi). \quad (4.18)$$

Lemma 4.4. *For any $\delta \in (0, 1]$, as $t \rightarrow \infty$ and $x/\sqrt{t} \rightarrow \infty$*

$$E_0[\text{vol}_d(S_A(\delta t)) \mid B_t = \mathbf{x}] = J(\delta t) + o(x \vee t). \quad (4.19)$$

Proof. The proof is given only for the case $\delta = 1$, the case $\delta < 1$ being dealt with in the same way. By (4.4) $E_0[\text{vol}_d(S_A(t)) \mid B_t = \mathbf{x}]$ is expressed as

$$\frac{1}{p_t(x)} \int_{\mathbf{z} \in \Omega_A} |d\mathbf{z}| \int_0^t ds \int_{\xi \in \partial A} H_A(\mathbf{z}, s; d\xi) p_{t-s}(\mathbf{x} - \mathbf{z} + \xi) + \text{vol}_d(A). \quad (4.20)$$

Step 1. We first consider this integral restricted to the region

$$D_t = \left\{ (\mathbf{z}, s) : \left| \mathbf{z} - \frac{s}{t} \mathbf{x} \right| < \sqrt{8T \lg T}, \quad 4 < s < t - 4, \mathbf{z} \notin A \right\}, \quad (4.21)$$

where $T = s(t-s)/t$. Observing that

$$\frac{\mathbf{x} - \mathbf{z}}{t-s} = \frac{\mathbf{x}}{t} - \frac{\mathbf{y}}{t-s} \quad \text{where} \quad \mathbf{y} := \mathbf{z} - \frac{s}{t} \mathbf{x}, \quad (4.22)$$

we obtain within D_t

$$\frac{p_{t-s}(\mathbf{x} - \mathbf{z} + \xi)}{p_{t-s}(\mathbf{x} - \mathbf{z})} = \exp \left[\frac{-(\mathbf{x} - \mathbf{z}) \cdot \xi - \frac{1}{2} |\xi|^2}{t-s} \right] = \exp \left[-\frac{\mathbf{x} \cdot \xi}{t} + O\left(\frac{\sqrt{T \lg T}}{t-s}\right) \right]. \quad (4.23)$$

Since $\sqrt{T \lg T}/(t-s) = \sqrt{\frac{s}{t(t-s)} \lg \frac{s(t-s)}{t}}$ is bounded for $s < t-4$ and approaches zero as $t-s \rightarrow \infty$, this results in the formula of the lemma, if the contributions from the complement of D_t to the integral (4.21) and the integral defining $J(t)$ are suitably evaluated, of which the former is dealt with in Step 2 and the latter in Step 3. In the case $x/t < M$ the evaluations may be readily performed in the same way as for that of the integral on $D_>$ in the proof of Lemma 4.2.

Step 2. Assume $x/t > 1$, although the following argument can apply in general. Put $\hat{D}_t = (\mathbf{R}^d \setminus A) \times [0, t] \setminus D_t$. Recollecting the proof of Lemma 4.1, we consider the inclusion

$$\{\mathbf{z} : \exists s, s = \sigma_{-A+\mathbf{z}} \text{ and } (\mathbf{z}, s) \in \hat{D}_t\} \subset \{\mathbf{z} : \exists s, \mathbf{z} - B_s \in \partial A \text{ and } (\mathbf{z}, s) \in \hat{D}_t\}$$

in place of (4.5). These sets depend on the Brownian path and are not empty only if it belongs to

$$\mathcal{E} := \{\exists s, (B_s + \xi, s) \in \hat{D}_t \text{ for some } \xi \in \partial A\}.$$

We observe that the contribution from $\hat{D}_t$ to the first term on the right-hand side of (4.20) is dominated by

$$\begin{aligned} & E[\text{vol}_d(\{\mathbf{z} : \exists s, s = \sigma_{-A+\mathbf{z}} \text{ and } (\mathbf{z}, s) \in \hat{D}_t\}) \mid B_t = \mathbf{x}] \\ & \leq E[\text{vol}_d(S_A(t)); \mathcal{E} \mid B_t = \mathbf{x}]. \end{aligned}$$

In view of (ii) of Lemma 4.3 (applied with $a = R_A$) and Corollary 4.1 (with ε in place of δ) an application of Schwarz inequality deduces that for each $\varepsilon > 0$ the last conditional expectation is at most $\text{vol}_{d-1}(\text{pr}_{\mathbf{e}}K)$ times

$$\left(P \left[\exists s \in [\varepsilon t, (1 - \varepsilon)t], (B_s + \xi, s) \in \hat{D}_t \text{ for some } \xi \in \partial A \mid B_t = \mathbf{x} \right] \right)^{1/2} x + \varepsilon x.$$

We claim that for each $\varepsilon > 0$ the conditional probability above tends to zero. For the proof we may disregard the dependence on ξ of the event under P . Now by scaling property of Brownian motion we infer that

$$\begin{aligned} & P[\exists s \in [\varepsilon t, (1 - \varepsilon)t], (B_s, s) \in \hat{D}_t \mid B_t = \mathbf{x}] \\ &= P[\exists u \in [\varepsilon, (1 - \varepsilon)], |B_u - u\mathbf{x}/\sqrt{t}| \geq \sqrt{8(u(1 - u) \lg T)} \mid B_1 = \mathbf{x}/\sqrt{t}] \\ &= P[\exists u \in [\varepsilon, (1 - \varepsilon)], |B_u| \geq \sqrt{8(u(1 - u) \lg[tu(1 - u)])} \mid B_1 = 0] \\ &\rightarrow 0 \end{aligned}$$

as $t \rightarrow \infty$, verifying the claim, which entails that

$$E_0[\text{vol}_d(S_A(t)) \mid B_t = \mathbf{x}] \leq J(t) + o(x).$$

Step 3. We must still show that the contribution from $\hat{D}_t \cap \mathbf{R}^d \times [4, t - 4]$ to the integral defining $J(t)$ is negligible, namely

$$\frac{1}{p_t(x)} \int_4^{t-4} ds \int_{|\mathbf{z}| \geq \sqrt{8T \lg T}} p_{t-s}(\mathbf{x} - \mathbf{z}) |d\mathbf{z}| ds \int_{\partial A} e^{-\mathbf{x} \cdot \xi / t} H_A(\mathbf{z}, s; d\xi) = o(x) \quad (4.24)$$

(see (4.18)). We may suppose that $v \rightarrow \infty$. Putting $R = 3R_A$, we take $\sqrt{8T(Rv \vee \lg T)}$ in place of $\sqrt{8T \lg T}$ in the definition of D_t and denote the resulting region by D'_x :

$$D'_x = \left\{ (\mathbf{z}, s) : \left| \mathbf{z} - \frac{s}{t} \mathbf{x} \right| < \sqrt{8(Rv \vee \lg T)T}, 4 < s < t - 4, \mathbf{z} \notin A \right\},$$

and its complement by $\hat{D}'_x$.

We first notify that the triple integral in (4.24) over $(\hat{D}'_x \cap \mathbf{R}^d \times [4, t - 4]) \times \partial A$ is evaluated to be $o(e^{-Rv} p_t(x) t)$. Indeed, using Lemma 3.5 and Theorem 2.1 in turn, we see $\int_{\partial A} e^{-\mathbf{x} \cdot \xi / t} H_A(\mathbf{z}, s; d\xi) \leq e^{R_A v} q_A(\mathbf{z}, s) \leq C \beta_A e^{Rv} p_s(\mathbf{z})$, and by using the identity (4.8) what is stated above is readily verified.

If $v < \lg(t \vee 2)$ is bounded, then the integral at issue restricted to $(D'_x \setminus D_t) \cap \mathbf{R}^d \times [4, t/2]$ is negligible owing to what is verified in Step 2, for the O term in (4.23), which is now $O(\sqrt{T(v \vee \lg T)}/(t - s))$, is bounded for $s < t/2$ so that we have the same integral as in (4.20). Therefore we have only to evaluate the integral in (4.24) in the situation when

$$\text{either (i) } v \geq \lg(t \vee 2) \quad \text{or (ii) } s < t/2 \text{ if } v \geq \lg(t \vee 2). \quad (4.25)$$

The integral defining $J(t)$ restricted to $\mathbf{R}^d \times [4, t - 4] \setminus D'_x$ is negligible as notified above and we compute the integral over $D'_x \setminus D_t$. Putting $\mathbf{v} = \mathbf{x}/t$ and

$$\varphi(\xi, u) = \int_{\partial A} e^{-\mathbf{v} \cdot \xi_1} H_A(\xi, u; d\xi_1),$$

we write the inner integral in (4.24) in the form

$$\int_{\partial A} e^{-\mathbf{x} \cdot \xi_1/t} H_A(\mathbf{z}, s; d\xi_1) = \int_0^s du \int_{\partial U(R)} \varphi(\xi, u) H_{U(R)}(\mathbf{z}, s-u; d\xi).$$

Let $I_{[\alpha, \beta]}$ denote the last repeated integral restricted to $[\alpha, \beta] \times \partial U(R)$. The contribution coming from the small interval $[0, M/v]$ is dominant and problematic. We first ascertain that the other part is negligible. If $(\mathbf{z}, s) \in D'_x \setminus D_t$ so that under (4.25)

$$\left| \frac{\mathbf{z}}{s} - \frac{\mathbf{x}}{t} \right| \leq \sqrt{4(Rv \vee \lg T)(t-s)/st} = O(\sqrt{v}), \quad (4.26)$$

then for t large enough,

$$I_{[M/v, s]} \leq e^{R_A v} \int_{M/v}^s q(z, s-u; R) du \leq c e^{2Rv} v^{\frac{1}{2}-\nu} p_{s-M/v}(z)$$

and $z^2/(s-M/v) - z^2/s > Mz^2/s^2 v > \frac{1}{2}vM$ (where $z = |\mathbf{z}|$); hence

$$I_{[M/v, s]} \leq p_s(z) e^{2Rv} e^{-\frac{1}{5}vM},$$

from which we readily infer that the part $I_{[M/v, s]}$ is negligible if $M > 10R$.

To deal with $I_{[0, M/v]}$ we are to apply Theorem 2.5. Define the measure $\mu_{s-u, \mathbf{z}}(d\xi)$ on $\partial U(R)$ via the equation

$$H_{U(R)}(\mathbf{z}, s-u; d\xi) = \omega_{d-1} R^{2\nu} v e^{\xi \cdot \mathbf{z}/(s-u)} p_{s-u}(z) \mu_{s-u, \mathbf{z}}(d\xi)$$

Recall the variable $\mathbf{y}$ that is introduced in (4.22) so that $\mathbf{z} = s\mathbf{v} + \mathbf{y}$. Let $0 \leq u \leq M/v$ and (4.25) be satisfied. Then, on writing $\mathbf{z} = s\mathbf{v} - u\mathbf{v} + \mathbf{y} - u\mathbf{y}/s + u\mathbf{z}/st$,

$$\frac{\mathbf{z}}{s-u} = \frac{\mathbf{x}}{t} + \frac{\mathbf{y}}{s} + O(1/s),$$

and, owing to (4.26), $|\mathbf{z}/z - \mathbf{x}/x| = O(1/\sqrt{v})$ and from Theorem 2.5 we infer that

$$\mu_{s-u, \mathbf{z}}(d\xi) \implies \frac{(\mathbf{x} \cdot \xi)}{x} m_R(d\xi) \mathbf{1}(|\theta| > \pi/2)$$

where θ is the angle formed by $\mathbf{x}$ and $\xi \in \partial U(R)$ so that $\cos \theta = \mathbf{x} \cdot \xi / xR$. (In the formula above the limit measure may depend on $\mathbf{e} = \mathbf{x}/x$ which is not fixed, but the convergence makes sense because of the uniformity of it for $\mathbf{e}$.) Thus the integral $I_{[0, M/v]}$ is essentially given by

$$[\omega_{d-1} R^{2\nu} v] \int_0^{M/v} p_{s-u}(z) du \int_{\partial U(R)} \varphi(\xi, u) e^{Rv \cos \theta} e^{\xi \cdot \mathbf{y}/s} \cos \theta \mathbf{1}(|\theta| > \pi/2) m_R(d\xi).$$

Noting that $\sup_{\xi, \xi' \in \partial U(R)} \varphi(\xi', u)/\varphi(\xi, u) \leq e^{2vR_A}$ and $|\mathbf{y}|/s = O(\sqrt{v})$ we compare the contributions from $\cos \theta > 9/10$ and $\cos \theta < 1/10$ to see that the integral w.r.t. ξ may be restricted to $\cos \theta > 1/10$. We then consider the integration w.r.t. $\mathbf{y}$ and compare the contribution from $y = |\mathbf{y}| \leq \sqrt{8T \lg T}$ and that from $\sqrt{8T \lg T} < y \leq \sqrt{8Rv}$. To this end put $T' = (s-u)(t-s)/(t-u)$ so that

$$p_{s-u}(\mathbf{z}) p_{t-s}(\mathbf{x} - \mathbf{z}) = p_{t-u}(\mathbf{x}) p_{T'}\left(\mathbf{z} - \frac{s-u}{t-s} \mathbf{x}\right).$$

Then, observing $T' \sim T$, $s > T'$ and

$$\mathbf{z} - \frac{s-u}{t-s}\mathbf{x} = \mathbf{y} + \frac{u(t-s)}{t(t-u)}\mathbf{x} = \mathbf{y} + b(\mathbf{y})$$

with $b(\mathbf{y}) = b_{s,t,u,\mathbf{x}}(\mathbf{y}) = O(1)$, we infer that the ratio of the integral

$$\int_{\sqrt{8T \lg T} < |\mathbf{y}| \leq \sqrt{8RTv}} e^{\xi \cdot \mathbf{y}/s} p_{T'}(\mathbf{y} + b(\mathbf{y})) |d\mathbf{y}|$$

to the same integral but over $y \leq \sqrt{8RT \lg T}$ tends to zero, which shows the required estimate of (4.24). This completes the proof of Lemma 4.4. $\square$.

Lemma 4.5. (i) For any Borel set $E \subset \partial A$ and $0 < s < t$,

$$H_A(\mathbf{x}, t; E) \leq \int_{\mathbf{z} \in \Omega_A} H_A(\mathbf{z}, s; E) p_{t-s}^{(d)}(\mathbf{z} - \mathbf{x}) |d\mathbf{z}|. \quad (4.27)$$

(ii) For each $\varepsilon \in (0, 1)$ there exists a constant M such that if $|\mathbf{x}|/\sqrt{t} > M$ and $t > M$, then for all $s \in [\varepsilon t, t]$ and $E \subset \partial A$,

$$\begin{aligned} H_A(\mathbf{x}, t; E) &\geq (1 - \varepsilon) \int_{\mathbf{z} \in \Omega_A} H_A(\mathbf{z}, s; E) p_{t-s}^{(d)}(\mathbf{z} - \mathbf{x}) |d\mathbf{z}| \\ &\quad - C \beta_A p_t^{(d)}(x) e^{-\varepsilon x^2/5t}, \end{aligned} \quad (4.28)$$

where β_A equals 1 or $\text{Cap}(A)$ according as $d = 2$ or $d \geq 3$ and C is a universal constant.

Proof. Let $0 < s < t$. By the strong Markov property of Brownian motion

$$H_A(\mathbf{x}, t; E) = \int_{\mathbf{z} \in \Omega_A} H_A(\mathbf{z}, s; E) P_{\mathbf{x}}[B_{t-s} \in d\mathbf{z}, \sigma_A > t - s];$$

in particular we have the upper bound (4.27).

As for the lower bound (4.28) let $s \geq \varepsilon t$ and first consider the contribution from $D_s := \{\mathbf{z} : |\mathbf{z} \cdot \mathbf{e}| < \frac{1}{3}sx/t\}$, $\mathbf{e} = \mathbf{x}/x$ to the integral in (4.27). To this end we simply replace E by ∂A , and apply Lemma 3.5 (or for $R_A x/t < 1$, Lemma 3.4) and Theorem 2.1 in turn to see that

$$H_A(\mathbf{z}, s; \partial A) \leq C \beta_A q(z, s; R) \leq C' \beta_A R^{2\nu} [(Rz/s) \vee 1]^{\frac{1}{2}-\nu} p_s^{(d)}(\mathbf{z}) e^{Rz/s}$$

($z = |\mathbf{z}|$). Writing $z_1 = \mathbf{z} \cdot \mathbf{e}$ and $y = |\mathbf{z} - z_1 \mathbf{e}|$, we deduce from the identity (4.8) that

$$p_s^{(d)}(\mathbf{z}) p_{t-s}^{(d)}(\mathbf{z} - \mathbf{x}) = p_t^{(d)}(x) p_T^{(d-1)}(y) p_T^{(1)}\left(z_1 - \frac{s}{t}x\right), \quad T = s(t-s)/t.$$

For simplicity let $R_A = R/2 = 1$. On substituting the obvious bounds

$$|z_1 - \frac{s}{t}x| \geq \frac{2}{3}sx/t \quad \text{and} \quad z \leq y + \frac{1}{3}sx/t$$

valid on D_s and dominating $[(2z/s) \vee 1]^{\frac{1}{2}-\nu}$ by $e^{z/s}$, the integral in (4.27) restricted on D_s is at most

$$C'' \beta_A e^{x/t} p_t(x) \left[\frac{sx}{t} p_T^{(1)}\left(\frac{2sx}{3t}\right) \right] \int_{\mathbf{R}^{d-1}} p_T^{(d-1)}(\mathbf{y}) e^{3|\mathbf{y}|/s} |d\mathbf{y}|. \quad (4.29)$$

A simple computation shows that the integral equals $e^{9T/2s^2} \leq e^2$ and the quantity in the square brackets is bounded by

$$\sqrt{\frac{x^2 s}{t(t-s)}} \exp \left\{ -\frac{2x^2 s}{9t(t-s)} \right\} \leq c \sqrt{\varepsilon x^2 / t} e^{-2\varepsilon x^2 / 9t},$$

where the monotonicity of $\alpha e^{-\alpha^2/2}$, $\alpha > 1$ as well as the assumption $s > \varepsilon t$ is applied for the last inequality. It is easy to see that (4.29) is at most the second term on the right-hand side of (4.28), provided $x > 20/\varepsilon$.

It now suffices to show that as $t \rightarrow \infty$ and $|\mathbf{x}|/\sqrt{t} \rightarrow \infty$

$$\frac{\int_{\mathbf{z} \notin D_s} H_A(\mathbf{z}, s; E) P_{\mathbf{x}}[B_{t-s} \in d\mathbf{z}, \sigma_A < t-s]}{\int_{\mathbf{z} \in \Omega_A} H_A(\mathbf{z}, s; E) p_{t-s}(\mathbf{z} - \mathbf{x}) |d\mathbf{z}|} \rightarrow 0. \quad (4.30)$$

Since

$$P_{\mathbf{x}}[B_{t-s} \in d\mathbf{z}, \sigma_A < t-s] / |d\mathbf{z}| = P_{\mathbf{x}}[\sigma_A < t-s | B_{t-s} = \mathbf{z}] p_{t-s}(\mathbf{z} - \mathbf{x}),$$

we now have only to show that uniformly for $\mathbf{z} \notin D_s$ and s ,

$$P_{\mathbf{x}}[\sigma_A < t-s | B_{t-s} = \mathbf{z}] \rightarrow 0. \quad (4.31)$$

On expressing the Brownian bridge by means of free Brownian motion and reversing the time we may write $P_{\mathbf{z}}[B_u + \frac{u}{t-s}(\mathbf{x} - B_{t-s}) \in A \text{ for some } u \in (0, t-s)]$ for the probability in (4.31), which, scaling the space and time variables, we therefore can dominate by

$$\sup_{\mathbf{y}: \mathbf{y} \cdot \mathbf{e} > sx/3t\sqrt{t-s}} P_{\mathbf{y}} \left[B_u - uB_1 + \frac{u\mathbf{x}}{\sqrt{t-s}} \in \frac{1}{\sqrt{t-s}} A \text{ for some } u \in (0, 1) \right].$$

Since $B_u - uB_1$ under $P_{\mathbf{y}}$ has the same law as $B_u - uB_1 + (1-u)\mathbf{y}$ under $P_{\mathbf{0}}$, and $sx/3t\sqrt{t-s} \rightarrow \infty$ so that $|(1-u)\mathbf{y} + u\mathbf{x}/\sqrt{t-s}| \rightarrow \infty$, the probability above obviously converges to zero. $\square$

Proof of Proposition 4.1. Perform the integration by $\mathbf{z}$ first in the triple integral defining $J(t) - J(\varepsilon t + 4)$ and substitute the inequality (4.28) and you find that

$$\begin{aligned} J(t) - J(\varepsilon t + 4) &= \frac{1}{p_t(x)} \int_{\mathbf{z} \in \Omega_A} |d\mathbf{z}| \int_{\varepsilon t}^{t-4} p_{t-s}(\mathbf{x} - \mathbf{z}) ds \int_{\partial A} e^{-\xi \cdot \mathbf{x}/t} H_A(\mathbf{z}, s; d\xi) \\ &\leq \int_{\xi \in \partial A} e^{-\xi \cdot \mathbf{x}/t} H_A(\mathbf{x}, t; d\xi) \frac{t}{p_t(x)} + tC\beta_A e^{-\varepsilon x^2/5t} \sup_{\xi \in A} e^{-\mathbf{x} \cdot \xi/t}. \end{aligned}$$

Now we apply (4.19) of Lemma 4.4 to see first that $J(\varepsilon t) \leq C\varepsilon(t \vee x)$ and then that the bound above gives the upper bound for the required equality. Use (4.27) instead of (4.28) to obtain the lower bound. $\square$

4.2.4. PROOFS OF THEOREMS 3.5 AND 4.3. Let K be a compact set and remember the measures $m_{K,\mathbf{e}}(d\xi)$ (defined in (3.35)) and

$$\mu_{t,\mathbf{x}}^K(d\xi) = \frac{e^{-v\mathbf{e} \cdot \xi} H_K(\mathbf{x}, t; d\xi)}{vp_t(x)} \quad (\mathbf{e} = \mathbf{x}/x, v = x/t).$$

According to Proposition 4.1 for any bounded Borel set A ,

$$E_{\mathbf{0}}[\text{vol}_d(S_A(t)) | B_t = \mathbf{x}] / x = \mu_{t,\mathbf{x}}^A(\partial A)(1 + o(1)). \quad (4.32)$$

Lemma 4.6. *The measure $\mu_{t,\mathbf{x}}^K|_{\langle K \rangle_{\mathbf{e}}}(d\xi)$ (restriction on $\langle K \rangle_{\mathbf{e}}$) is asymptotically dominated by $m_{K,\mathbf{e}}(d\xi)$ in the sense that for any Borel set $E \subset \langle K \rangle_{\mathbf{e}}$*

$$\limsup_{x/t \rightarrow \infty, t \rightarrow \infty} \left[\mu_{t,\mathbf{x}}^K(E) - m_{K,\mathbf{e}}(E) \right] \leq 0. \quad (4.33)$$

Proof. From the inclusion $E \subset \partial K$ it follows that $H_K(\mathbf{x}, t; d\xi) \leq H_E(\mathbf{x}, t; d\xi)$ for $d\xi \subset E$ since Brownian paths that hits $\partial K \setminus E$ before E may contribute to H_E but never to $H_{\partial K}|_E$. Hence

$$\mu_{t,\mathbf{x}}^K(E) \leq \mu_{t,\mathbf{x}}^E(E).$$

By (4.32) the right-hand side is asymptotically dominated by $E[\text{vol}_d(S_E(t))|B_t = \mathbf{x}]/x$, which in turn is asymptotically dominated by $\text{vol}_{d-1}(\text{pr}_{\mathbf{e}}(E)) = m_{K,\mathbf{e}}(E)$ owing to Lemma 4.3, showing the required bound. $\square$

Corollary 4.3. *Let $\mathbf{e} \in \partial U(1)$ and $\mathbf{x} = x\mathbf{e}$ and suppose that $\langle K \rangle_{\mathbf{e}}$ lies on a plane perpendicular to $\mathbf{e}$. Then as $x/t \rightarrow \infty$ and $t \rightarrow \infty$, $\mu_{t,\mathbf{x}}^K \implies m_{K,\mathbf{e}}$, in which the two sides of K are distinguished and the measure $m_{K,\mathbf{e}}$ is considered to be concentrated on the $+\mathbf{e}$ side of K .*

Proof. This corollary follows from the proof of Lemma 4.6 (either directly or with the help of Proposition 4.1 and Corollary 4.2). $\square$

Proof of Theorem 3.5. Let $V_\beta = \{\mathbf{z} \in \mathbf{R}^d : \mathbf{z} \cdot \mathbf{e} \geq -\beta\}$ for $\beta > 0$ and, choosing β so that $K \subset V_\beta$, put

$$L_\xi = \{\xi - s\mathbf{e} : s \geq 0\} \cap V_\beta \quad \text{for } \xi \in \langle K \rangle_{\mathbf{e}} \quad \text{and} \quad W = \cup_{\xi \in \langle K \rangle_{\mathbf{e}}} L_\xi.$$

Note that $K \subset W$, $\langle W \rangle_{\mathbf{e}} = \langle K \rangle_{\mathbf{e}}$ and W is compact. Let

$$D = W \cap \partial V_\beta \quad \text{and} \quad C = \partial W \setminus (\langle K \rangle_{\mathbf{e}} \cup D).$$

Then C, D and $\langle K \rangle_{\mathbf{e}}$ together make up the decomposition of ∂W .

Now consider the limit procedure as $t \rightarrow \infty, x/t \rightarrow \infty$. Since

$$\text{pr}_{\mathbf{e}} C = \text{dis-ct}_{\mathbf{e}}(K),$$

the assumption (3.37) implies $\text{vol}_{d-1}(\overline{\text{pr}_{\mathbf{e}} C}) = 0$; hence $E[S_C(t)|B_t = \mathbf{x}] = o(x)$ owing to Lemma 4.3, which in turn shows that $\mu_t^C(C) \rightarrow 0$. By Corollary 4.3 $\mu_{t,\mathbf{x}}^W(D) \rightarrow 0$. Thus the limit measure of μ_t^W must concentrate on $\langle K \rangle_{\mathbf{e}}$, hence is not larger than $m_{K,\mathbf{e}} (= m_{W,\mathbf{e}})$ in view of Lemma 4.6, but by Proposition 4.1 it cannot be strictly less than $m_{K,\mathbf{e}}$ since according to Corollaries 4.1 and 4.2

$$E[S_W(t) | B_t = \mathbf{x}] \sim m_{K,\mathbf{e}}(K)vt. \quad (4.34)$$

Finally we have the inequality $\mu_{t,\mathbf{x}}^W(d\xi) \leq \mu_{t,\mathbf{x}}^K(d\xi)$ for $d\xi \subset \langle K \rangle_{\mathbf{e}}$ since $K \subset W$. Applying Proposition 4.1 again (but this time to K) we obtain (3.38) as required. $\square$

Proof of Theorem 4.3. This is now clear from Theorem 3.5 and Proposition 4.1. $\square$

5 Brownian Motion with A Constant Drift $-v\mathbf{e}$

The Brownian bridge $P_0[\cdot | B_t = x\mathbf{e}]$ with $v := x/t$ kept away from zero may be comparable or similar to the process $B_s - sv\mathbf{e}$, $0 \leq s \leq t$ in significant respects and here we present the results for the latter that are readily derived from those given above for the bridge.

We fix a unit vector $\mathbf{e} \in \partial U(1)$ ($d \geq 2$). Given $v > 0$, we put $\mathbf{v} = v\mathbf{e}$ and label with the superscript $(\mathbf{v})$ the objects defined by means of $B_s - \mathbf{v}s$ in place of B_t ; in particular, $B_s^{(\mathbf{v})} := B_s - \mathbf{v}s$. Let A be bounded Borel and consider

$$H_A^{(\mathbf{v})}(\mathbf{x}, s; d\xi) = P_{\mathbf{x}}[B_{\sigma^{(\mathbf{v})}(A)}^{(\mathbf{v})} \in d\xi, \sigma_A^{(\mathbf{v})} \in ds]/ds.$$

We put $\gamma(\cdot) = -\mathbf{v}$ (constant function) and $Z(s) = e^{\int_0^s \gamma(B_u)dB_u - \frac{1}{2} \int_0^s \gamma^2(B_u)du}$. Then, for a bounded continuous function ψ on $[0, \infty) \times \partial A$,

$$\int_{[0, \infty) \times \partial A} \psi(u, \xi) H_A^{(\mathbf{v})}(\mathbf{x}, u; d\xi) du = E_{\mathbf{x}}[Z(\sigma_A) \psi(\sigma_A, B_{\sigma(A)})]$$

owing to the Cameron-Martin formula. Since

$$Z(\sigma(A)) = \exp \left\{ -\mathbf{v} \cdot B_{\sigma(A)} + \mathbf{v} \cdot \mathbf{x} - \frac{1}{2} v^2 \sigma_A \right\},$$

we obtain

$$H_A^{(\mathbf{v})}(\mathbf{x}, s; d\xi) = e^{\mathbf{v} \cdot \mathbf{x} - \frac{1}{2} v^2 s} e^{-\mathbf{v} \cdot \xi} H_A(\mathbf{x}, s; d\xi). \quad (5.1)$$

On writing $\mathbf{x} = \mathbf{v}t + \mathbf{y}$, this together with the first formula (vi) of Section 1 yields

$$(vi') \quad H_A^{(\mathbf{v})}(\mathbf{x}, t; d\xi) \sim \Lambda_{\nu}(R_A v) (2\pi t)^{-d/2} e^{\mathbf{v} \cdot \mathbf{y} - \mathbf{v} \cdot \xi} \lambda_A(\mathbf{x}/t; d\xi) \quad (d\xi \subset \partial A)$$

with $\lambda_A(\mathbf{v}; d\xi)$ defined in (4.14).

From the fact that the law of Brownian bridge does not depends on the strength of drift, we have

$$E_0[\text{vol}_d(S_A^{(\mathbf{v})}(t))] = \int_{\mathbf{R}^d} E_0[\text{vol}_d(S_A(t)) | B_t = \mathbf{x}] p_t^{(d)}(\mathbf{x} - \mathbf{v}t) d\mathbf{x}, \quad (5.2)$$

Noting that overwhelming contribution to the integral comes from the relatively small range $\mathbf{v}t + U(t^\alpha)$ with any α such that $1/2 < \alpha < 1$ we infer from the second formula of (vi) that as $t \rightarrow \infty$, uniformly for v in a bounded interval,

$$(vi'') \quad E_0[\text{vol}_d(S_A^{(\mathbf{v})}(t))] \sim \Lambda_{\nu}(R_A v) t \int_{\partial A} e^{-\mathbf{v} \cdot \xi} \lambda_A(\mathbf{v}, d\xi),$$

For the case of a disc/ball it follows from (1.9) that

$$E_0[\text{vol}_d(S_{U(a)}^{(\mathbf{v})}(t))] \sim a^{2\nu} \Lambda_{\nu}(av) t \int_{\partial U(1)} e^{-a\mathbf{v} \cdot \xi} g_{av}(\theta) m_1(d\xi)$$

with $g_a(\theta)$ given in (2.5) of Section 2. Here $\theta = \theta(\xi)$ denotes the angle ξ forms with $\mathbf{e}$.

Consider the case when $v \rightarrow \infty$ as well as $t \rightarrow \infty$. Let K be a compact set that satisfies the condition (3.37), i.e., $\text{vol}_{d-1}(\overline{\text{dis-ct}_{\mathbf{e}}(K)}) = 0$. Then, substituting $s = t + \tau$ (with $|\tau| \ll t$) into (5.1), and comparing the resulting formula with

$$p_{t+\tau}(vt) = p_t(vt) \exp \left\{ -\frac{1}{2} v^2 \tau \left(1 - \frac{\tau}{t} + \frac{\tau^2}{t^2} - \dots \right) \right\},$$

we infer from (vii) that uniformly for τ in a finite interval, as $v \rightarrow \infty$ with $v/t \rightarrow 0$ (so that the ratio $\mathbf{v} \cdot \xi/t$ as well as the third term in the exponent tends to zero)

$$(vii') \quad H_K^{(\mathbf{v})}(\mathbf{x}, t + \tau; d\xi) \sim v p_t(v\tau) m_{K, \mathbf{x}/x}(d\xi) \quad (d\xi \subset \partial K)$$

for $\mathbf{x} = \mathbf{v}t$, which may be broken into the two relations

$$q_K^{(\mathbf{v})}(\mathbf{x}, t + \tau) \sim v p_t(v\tau) \text{vol}_{d-1}(\text{pr}_{\mathbf{x}/x} K) \quad (5.3)$$

and

$$P_{\mathbf{x}}[B_t^{(\mathbf{v})} \in d\xi \mid \sigma_K^{(\mathbf{v})} = t + \tau] \sim \frac{m_{K, \mathbf{x}/x}(d\xi)}{\text{vol}_{d-1}(\text{pr}_{\mathbf{x}/x} K)}. \quad (5.4)$$

In these formulae, as is readily ascertained, we may take $\mathbf{x} = \mathbf{v}t + \mathbf{y}$ in place of $\mathbf{x} = \mathbf{v}t$, provided that $\mathbf{v} \cdot \mathbf{y} \rightarrow 0$ and $|\mathbf{y}|/t \rightarrow 0$.

Also from (viii) and (5.2) it follows that as $v \rightarrow \infty$ and $t \rightarrow \infty$

$$(viii') \quad E_0[\text{vol}_d(S_K^{(\mathbf{v})}(t))] \sim [\text{vol}_{d-1}(\text{pr}_{\mathbf{e}} K)] vt \quad (\mathbf{v} = v\mathbf{e}).$$

The formula (5.4) (at least with $\tau = 0$) (as well as (viii')) is intuitively comprehensible, but it (hence also (5.3)) is not true (for any τ) if $v/\sqrt{t}$ is bounded away from zero when the ratio $P_{\mathbf{x}}[\sigma_{\Delta_1}^{(\mathbf{v})} \in dt]/P_{\mathbf{x}}[\sigma_{\Delta_2}^{(\mathbf{v})} \in dt]$ does not approaches unity, where Δ_1 and Δ_2 are any (but distinct) two fixed hyperplane perpendicular to $\mathbf{e}$.

6 Appendix

The arguments presented in this section except the last subsection are made independently of those of preceding sections. Throughout this section A denotes a bounded Borel set of $\mathbf{R}^d$ and put $R_A = \sup\{|\mathbf{y}| : \mathbf{y} \in A\}$.

6.1 Harmonic measure of heat operator

Here we give a brief exposition of the well known fact that $H_A(\mathbf{x}, t; d\xi)dt$ is the lateral component of the caloric measure for the exterior of the cylinder with base A . Given $t > 0$, the space-time Brownian motion $Y_s = (B_s, t - s)$ ($0 \leq s \leq t$) under the law $P_{\mathbf{x}}$ is regulated by the heat operator $\frac{1}{2}\Delta - (\partial/\partial s)$. Let A^r designate the set of regular points of A and Ω_A the unbounded component of $\mathbf{R}^d \setminus A^r$ as in Introduction, put

$$D = \Omega_A \times (0, \infty) = \{(\mathbf{z}, t) : \mathbf{z} \in \Omega_A, t > 0\}$$

and consider the following ‘Dirichlet problem’ for the heat operator:

$$\begin{aligned} \left(\frac{1}{2}\Delta - \frac{\partial}{\partial t}\right)u &= 0 \quad \text{on } D \\ u &= \varphi \quad \text{on } \partial_{\text{reg}} D := A^r \times [0, \infty) \cup \Omega_A \times \{t = 0\}, \end{aligned} \quad (6.1)$$

where, φ is any bounded continuous function on $\partial_{\text{reg}} D$ and the boundary condition (6.1) is interpreted in a reasonable way. The caloric measure, $\mu_D(\mathbf{x}, t, d\xi ds)$ say, for D at a reference

point $(\mathbf{x}, t) \in D$ is defined as such a measure kernel that the above boundary value problem may be solved in the form

$$u(\mathbf{x}, t) = \int_{\partial_{\text{reg}} D} \varphi(\xi, s) \mu_D(\mathbf{x}, t; d\xi ds),$$

whereas the solution to the same problem is represented by the expectation:

$$u(\mathbf{x}, t) = E_{\mathbf{x}}[\varphi(B_{\sigma(A)}, t - \sigma_A); \sigma_A < t] + E_{\mathbf{x}}[\varphi(B_t, 0); \sigma_A > t].$$

The first expectation on the right-hand side above is expressed as an integral by the measure kernel $Q_A(\mathbf{x}, dt d\xi)$ given in (1.1). The function $u^A(\mathbf{x}, t) := P_{\mathbf{x}}[\sigma_A \leq t] = Q(\mathbf{x}, \partial A \times [0, t])$ satisfies the heat equation in the interior D° , hence its partial derivative $q_A(\mathbf{x}, t) = u_t^A(\mathbf{x}, t)$ is not only well-defined but smooth in D° as is u^A ; hence also the derivative $H_A(\mathbf{x}, t; d\xi) = Q_A(\mathbf{x}, dt d\xi)/dt$. This assures that the caloric measure μ_D restricted to the lateral part of $\partial_{\text{reg}} D$ is written as

$$\mu_D(\mathbf{x}, t; d\xi dt) \Big|_{A^r \times [0, \infty)} = H(\mathbf{x}, t - s; d\xi) ds,$$

thus providing the probabilistic expression of μ_D (cf. Hunt [10]), the one on the initial boundary $\Omega_A \times \{0\}$ being of course given by $P_{\mathbf{x}}[B_t \in d\xi, \sigma_A > t] \ (\xi \in \Omega_A)$.

6.2 Uniform estimates for $e_A(\mathbf{x})/P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]$

Here we discuss a part of classical potential theory in two dimensions related to the function $e_A(\mathbf{x})$ and thereby prove Lemma 3.3. Most of what are presented are known but some as given in Proposition 6.1 below do not seem to be found in the existing literature.

Let $d = 2$. Suppose the bounded Borel set A to be non-polar. Hunt [10] (Section 5.5) defines $e_A(\mathbf{x})$ by

$$e_A(\mathbf{x}) = \pi g_{\Omega_A}(\mathbf{x}, \mathbf{y}) + \lg |\mathbf{x} - \mathbf{y}| - E_{\mathbf{x}}[\lg |B_{\sigma(A)} - \mathbf{y}|] \quad (6.2)$$

(H is written for e_A/π in [10]). Here g_{Ω_A} stands for the Green function for the domain Ω_A :

$$g_{\Omega_A}(\mathbf{x}, \mathbf{y}) = \int_0^\infty \frac{P_{\mathbf{x}}[B_t \in d\mathbf{y}, \sigma_A > t]}{|d\mathbf{y}|} dt \quad (\mathbf{x}, \mathbf{y} \in \Omega_A),$$

where $|\cdot|$ designates the Lebesgue measure on $\mathbf{R}^2$; $e_A(\mathbf{x})$ is defined for all $\mathbf{x} \in \mathbf{R}^2$ and independent of $\mathbf{y} \in \Omega_A$ (this fact is seen from the arguments developed for (6.5) below). It follows that

$$e_A(\mathbf{x}) = \pi \lim_{|\mathbf{y}| \rightarrow \infty} g_{\Omega_A}(\mathbf{x}, \mathbf{y}), \quad (6.3)$$

in particular e_A is harmonic in Ω_A . Lemma 3.3 readily follows from the following by using $(1+x)^{-1} > 1-x, x > 0$. Recall $R_A = \sup\{|\mathbf{y}| : \mathbf{y} \in A^r\}$.

Proposition 6.1. *For $r > R_A$ and $\mathbf{x} \in \Omega_A \cap U(r)$,*

$$P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] = \frac{e_A(\mathbf{x})}{\lg(r/R_A)} \left(1 + \frac{m_{R_A}(e_A)}{\lg(r/R_A)} (1 + \delta) \right)^{-1} \quad (6.4)$$

with $-2(R_A^{-1}r + 1)^{-1} \leq \delta \leq 2(R_A^{-1}r - 1)^{-1}$. Here $m_R(e_A) = \int_{\partial U(R)} e_A(\xi) m_R(d\xi)$.

In below we shall derive from (6.2) several formulae that relate $P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]$ and e_A ; Proposition 6.1 will be among them (see Corollary 6.1).

We begin with proving that for each $R > R_A$, uniformly for $\mathbf{x} \in U(R) \cap \Omega_A$

$$(\lg r)P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \rightarrow e_A(\mathbf{x}) \quad \text{as } r \rightarrow \infty, \quad (6.5)$$

which result is proved in [20]. We follow lines of [20]. Put $\Omega_r = U(r) \cap \Omega_A$ and let $\tau(\Omega_r)$ denote the first exit time from Ω_r . The proof is based on the identity

$$0 = \pi g_{\Omega_r}(\mathbf{x}, \mathbf{y}) + \lg |\mathbf{x} - \mathbf{y}| - E_{\mathbf{x}}[\lg |B_{\tau(\Omega_r)} - \mathbf{y}|] \quad (\mathbf{x}, \mathbf{y} \in \Omega_r, \mathbf{x} \neq \mathbf{y}). \quad (6.6)$$

We are to break the last expectations in (6.2) and (6.6) according as $\sigma_{\partial U(r)}$ is larger or smaller than σ_A . On noting that $\lg |B_{\sigma_A} - \mathbf{y}|$ agrees with $\lg |B_{\tau(\Omega_r)} - \mathbf{y}|$ a.s. on the event $\sigma_A < \sigma_{\partial U(r)}$ we observe

$$\begin{aligned} & E_{\mathbf{x}}[\lg |B_{\tau(\Omega_r)} - \mathbf{y}|] - E_{\mathbf{x}}[\lg |B_{\sigma(A)} - \mathbf{y}|] - (\lg r)P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \\ &= E_{\mathbf{x}}[\lg |B_{\tau(\Omega_r)} - \mathbf{y}| - \lg r; \sigma_{\partial U(r)} < \sigma_A] - E_{\mathbf{x}}[\lg |B_{\sigma(A)} - \mathbf{y}|; \sigma_{\partial U(r)} < \sigma_A], \end{aligned} \quad (6.7)$$

of which each of the expectations on the right-hand side approaches zero as $r \rightarrow \infty$. Now we consider the difference $e_A(\mathbf{x}) - (\lg r)P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]$. Substituting into it the expression of (6.2) and subtracting the right-hand side of (6.6), we find

$$\begin{aligned} & e_A(\mathbf{x}) - (\lg r)P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \\ &= \pi[g_{\Omega_A}(\mathbf{x}, \mathbf{y}) - g_{\Omega_r}(\mathbf{x}, \mathbf{y})] + \text{RHS of (6.7)} \end{aligned} \quad (6.8)$$

and readily conclude (6.5).

We bring in the function

$$\bar{e}_{A,R}(\mathbf{x}) = E_{\mathbf{x}}[e_A(B_{\sigma(U(R))})] \quad (x \geq R \geq R_A).$$

Then, the formula (6.5) entails

$$e_A(\mathbf{x}) = \lg(x/R) + \bar{e}_{A,R}(\mathbf{x}) \quad \text{whenever } x \geq R \geq R_A. \quad (6.9)$$

Indeed, for $\mathbf{x}$ with $R < x < r$,

$$\begin{aligned} & P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \\ &= P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_{\partial U(R)}] \\ &\quad + \int_{\partial U(R)} P_{\mathbf{x}}[\sigma_{\partial U(R)} < \sigma_{\partial U(r)}, B_{\sigma(U(R))} \in d\xi] P_{\xi}[\sigma_{\partial U(r)} < \sigma_A] \end{aligned} \quad (6.10)$$

and, noting that the first term on the right-hand side equals $\lg(x/R)/\lg(r/R)$, we multiply $\lg r$, let $r \rightarrow \infty$ and apply (6.5) to obtain (6.9).

Lemma 6.1. *Let $r > R \geq R_A$ and $R \leq x < r$. Then*

$$e_A(\mathbf{x}) = P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \left(\lg(r/R) + E_{\mathbf{x}}[\bar{e}_{A,R}(B_{\partial U(r)}) \mid \sigma_{\partial U(r)} < \sigma_A] \right). \quad (6.11)$$

Proof. Take $r^* > r$ and apply the strong Markov property to see that

$$P_{\mathbf{x}}[\sigma_{\partial U(r^*)} < \sigma_A] = P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \int_{\partial U(r)} P_{\xi}[\sigma_{\partial U(r^*)} < \sigma_A] \mu_{r,\mathbf{x}}(d\xi), \quad (6.12)$$

where

$$\mu_{r,\mathbf{x}}(d\xi) = P_{\mathbf{x}}[B_{\sigma(\partial U(r))} \in d\xi \mid \sigma_{\partial U(r)} < \sigma_A].$$

Then multiply the both sides by $\lg r^*$, let $r^* \rightarrow \infty$ and apply first the formula (6.5) with r^* in place of r , and then (6.9) with r in place of R to $e_A(\xi)$ that comes up on the right side under the integral sign, and one then finds the identity of the lemma. $\square$

By using an explicit form of the Poisson kernel for $\Omega_{U(R)} = \{\mathbf{z} \in \mathbf{R}^d : |\mathbf{z}| > R\}$, we have for $\mathbf{y} \in \partial U(r)$

$$\frac{r-R}{r+R} \leq \frac{P_{\mathbf{y}}[B_{\sigma(U(R))} \in d\xi]}{m_R(d\xi)} \leq \frac{r+R}{r-R} \quad (r > R),$$

entailing $m_R(e_A) \frac{r-R}{r+R} \leq \bar{e}_{A,R}(\mathbf{y}) \leq \frac{r+R}{r-R} m_R(e_A)$, which combined with Lemma 6.1 yields

Corollary 6.1. For $r > R_A$ and $\mathbf{x} \in \Omega_A \cap U(r)$,

$$m_{R_A}(e_A) \frac{r-R_A}{r+R_A} \leq \frac{e_A(\mathbf{x})}{P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]} - \lg \left(\frac{r}{R_A} \right) \leq m_{R_A}(e_A) \frac{r+R_A}{r-R_A}.$$

Rearranging this formula leads to Proposition 6.1.

The identity (6.9) shows

$$\lim_{x \rightarrow \infty} (e_A(\mathbf{x}) - \lg x) = -\lg R + m_R(e_A). \quad (6.13)$$

Since the left-hand side is independent of R , the right-hand side must be a constant depending only on A , which, known as *Robin's constant* associated with A , we denote by $V(A)$:

$$V(A) = -\lg R + m_R(e_A). \quad (6.14)$$

Since g_{Ω_A} is symmetric, letting $x \rightarrow \infty$ in (6.2) we find another (rather classical) representation

$$V(A) = e_A(\mathbf{y}) - \int_{\partial A} \lg |\xi - \mathbf{y}| H_A^\infty(d\xi), \quad \mathbf{y} \in \Omega_A. \quad (6.15)$$

Using the formula (6.15) instead of (6.9) in the very last step of the proof of Lemma 6.1, we deduce from (6.12) that

$$e_A(\mathbf{x}) = P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] \left[V(A) + \int_{\partial U(r)} \mu_{r,\mathbf{x}}(d\xi') \int_{\partial A} \lg |\xi - \xi'| H_A^\infty(d\xi) \right].$$

The repeated integral in the large square brackets may be written as $\lg r + \delta(r)$ with $|\delta(r)| \leq |\lg(1 - R_A/r)|$. Thus

$$|e_A(\mathbf{x})/P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A] - V(A) - \lg r| \leq -\lg(1 - R_A/r),$$

which in terms of the *logarithmic capacity* defined by

$$\text{lcap}(A) = e^{-V(A)} \quad (6.16)$$

(normalized so that $\text{lcap}(U(a)) = a$) may be expressed as in the following

Proposition 6.2. For $r > R_A$ and for $\mathbf{x} \in \Omega_A \cap U(r)$

$$\left| \frac{e_A(\mathbf{x})}{P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]} - \lg \left(\frac{r}{\text{lcap}(A)} \right) \right| \leq -\lg \left(1 - \frac{R_A}{r} \right). \quad (6.17)$$

By virtue of (6.14) the twin inequalities of Corollary 6.1 may be written as

$$-\frac{2m_{R_A}(e_A)}{r/R_A + 1} \leq \frac{e_A(\mathbf{x})}{P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]} - \lg \left(\frac{r}{\text{lcap}(A)} \right) \leq \frac{2m_{R_A}(e_A)}{r/R_A - 1},$$

which combined with Proposition 6.2 yields

Corollary 6.2. If $x > 2R_A$, then for some universal constant C ,

$$\left| \frac{e_A(\mathbf{x})}{P_{\mathbf{x}}[\sigma_{\partial U(r)} < \sigma_A]} - \lg \left(\frac{r}{\text{lcap}(A)} \right) \right| \leq C[1 \wedge m_{R_A}(e_A)] \frac{R_A}{r}.$$

6.3 An upper bound for q_A ($d \geq 3$)

Let $d \geq 3$ and let $\kappa_d, \kappa'_d, \kappa''_d$ etc. designate constants that depends only on d , whose precise values are not important to the present purpose and may vary from line to line.

In this subsection we prove

Proposition 6.3. For each $M \geq 1$, there exists a constant $\kappa_{d,M}$ (depending only on M and d) such that for all $t \geq R_A^2$ and $\mathbf{x} \in \Omega_A$ with $x/t < M/R_A$,

$$q_A(\mathbf{x}, t) \leq \kappa_{M,d} \text{Cap}(A) \Lambda_\nu(R_A x/t) p_t^{(d)}(x).$$

($\text{Cap}(A)$ is the Newtonian capacity defined in Section 3.1.1.)

The constant $\kappa_{M,d}$ in Proposition 6.3 may actually depend on M , the factor $\text{Cap}(A)$ being possibly too small for the inequality to be valid with a constant independent of M . This is caused by the concentration of the measure $H_{U(R_A)}(\mathbf{x}, t; d\xi)$ at $R_A \mathbf{x}/x$ as $x/t \rightarrow \infty$. (Compare with the result for $d = 2$ given in Proposition 6.4.)

The proof consists of several lemmas.

Lemma 6.2. Put $\eta_{\mathbf{x}} = \eta_{A,\mathbf{x}} = \text{dist}(\mathbf{x}, A^c)$. Then for all $\mathbf{x} \in \Omega_A$ and $t \geq 0$,

$$P_{\mathbf{x}}[t < \sigma_A < \infty] \leq \frac{\kappa_d}{(t \vee \eta_{\mathbf{x}}^2)^\nu} \text{Cap}(A).$$

Proof. Put $\mu(d\xi) = \mu_{t,\mathbf{x}}(d\xi) = P_{\mathbf{x}}[B_{\sigma(A)} \in d\xi, t < \sigma_A < \infty]$. If φ is a positive Borel function,

$$\begin{aligned} & \int \mu(d\xi) \int \varphi(\mathbf{z}) G^{(d)}(\xi - \mathbf{z}) |d\mathbf{z}| \\ &= E_{\mathbf{x}} \left[\int_0^\infty E_{B_{\sigma(A)}}[\varphi(B_s)] ds; t < \sigma_A < \infty \right] \\ &= E_{\mathbf{x}} \left[\int_{\sigma(A)}^\infty \varphi(B_s) ds; t < \sigma_A < \infty \right] \\ &\leq E_{\mathbf{x}} \left[\int_t^\infty \varphi(B_s) ds \right] = \int_t^\infty ds \int p_s^{(d)}(\mathbf{x} - \mathbf{z}) \varphi(\mathbf{z}) |d\mathbf{z}|. \end{aligned}$$

Note that the potential of μ is maximized on the set $\bar{A}$. Then, we infer from the inequality above that

$$\begin{aligned} \int G^{(d)}(\xi - \mathbf{z})\mu(d\xi) &\leq \sup_{\mathbf{z} \in \bar{A}} \int_t^\infty p_s^{(d)}(\mathbf{z} - \mathbf{x})ds \\ &\leq \int_t^\infty p_s^{(d)}(\eta_{\mathbf{x}})ds = \frac{1}{(2\pi)^{d/2}\eta_{\mathbf{x}}^{d-2}} \int_0^{\eta_{\mathbf{x}}/\sqrt{t}} u^{d-3} e^{-u^2/2} du. \end{aligned}$$

The right most member is at most $\kappa'_d(\sqrt{t} \vee \eta_{\mathbf{x}})^{-(d-2)} \kappa'_d[t \vee \eta_{\mathbf{x}}^2]^{-\nu}$ and the left most member is at least the ratio of the total measure of μ to $\text{Cap}(A)G^{(d)}(1)$, hence we have the inequality of the lemma with $\kappa_d = \kappa'_d/G^{(d)}(1)$. $\square$

Lemma 6.3. *Let $\eta_{\mathbf{x}} = \text{dist}(\mathbf{x}, A^c)$ as above. Then, (i) for all $\mathbf{x} \in \Omega_A$ and $t \geq 1$,*

$$q_A(\mathbf{x}, t) \leq \kappa_d(t \vee \eta_{\mathbf{x}}^2)^{-\nu} \text{Cap}(A);$$

and (ii) for all $t > 0$ and $\mathbf{x} \in \Omega_A$ with $\eta_{\mathbf{x}} \geq 1$, $q_A(\mathbf{x}, t) \leq \kappa_d \text{Cap}(A)$.

Proof. Let $t \geq 1$. Since then $\inf_{\mathbf{y} \in U(1)}(t \vee \eta_{\mathbf{x}+\mathbf{y}}^2) \geq \frac{1}{4}(t \vee \eta_{\mathbf{x}}^2)$, by the preceding lemma we have

$$\int_{t/3}^{2t/3} q_A(\mathbf{x} + \mathbf{y}, s)ds \leq \kappa'_d(t \vee \eta_{\mathbf{x}}^2)^{-\nu} \text{Cap}(A) \quad \text{for } \mathbf{y} \in U(1),$$

so that there exists $s^* \in [\frac{1}{3}t, \frac{2}{3}t]$ such that

$$\int_{U(1)} q_A(\mathbf{x} + \mathbf{y}, s^*)|d\mathbf{y}| \leq 3\kappa'_d t^{-1}(t \vee \eta_{\mathbf{x}}^2)^{-\nu} \text{Cap}(A). \quad (6.18)$$

Here we set $q_A(\mathbf{z}, \cdot) = 0$ for $\mathbf{z} \notin \Omega_A$. Denote by $\tau = \tau_{\mathbf{x}}$ the first exit time from the ball $\mathbf{x} + U(1)$. Then by strong Markov property

$$\begin{aligned} q_A(\mathbf{x}, t) &= \int_0^{t-s^*} \int_{\partial U(1)} P_{\mathbf{x}}[\sigma_A > s, \tau \in ds, B_\tau - \mathbf{x} \in d\xi] q_A(\mathbf{x} + \xi, t-s) \\ &\quad + \int_{U(1)} P_{\mathbf{x}}[B_{t-s^*} - \mathbf{x} \in d\mathbf{y}, \tau \wedge \sigma_A > t-s^*] q_A(\mathbf{x} + \mathbf{y}, s^*). \end{aligned} \quad (6.19)$$

Using the bound

$$P_{\mathbf{x}}[\sigma_A > s, \tau \in ds, B_\tau - \mathbf{x} \in d\xi] \leq \kappa''_d m_1(d\xi)ds, \quad (6.20)$$

together with Lemma 6.2 we infer that the first term (i.e., the repeated integral) is dominated by $4\kappa_d(t \vee \eta_{\mathbf{x}}^2)^{-\nu} \text{Cap}(A)$. As for the second term we see that for some universal constant $\lambda > 0$, $P_{\mathbf{x}}[B_{t-s^*} - \mathbf{x} \in d\mathbf{y}, \tau \wedge \sigma_A > t-s^*]/|d\mathbf{y}| \leq Ce^{-\lambda t}$, hence the bound (6.18) yields an estimate enough for the required estimate of (i).

If $\eta_{\mathbf{x}} \geq 1$ and $f(s) = P_{\mathbf{x}}[\tau_{\mathbf{x}} \in ds]/ds$ with the same $\tau_{\mathbf{x}}$ as above, then for $t < 1$, $f(1-s) \geq Cf(t-s)$ for some C so that

$$q_A(\mathbf{x}, 1) \geq \int_0^t f(1-s)ds \int q_A(\mathbf{x} + \xi, s)m_1(d\xi) \geq Cq_A(\mathbf{x}, t).$$

Hence (ii) follows from (i). $\square$

It is noted that in view of the scaling relations given in (1.11) the bound (ii) of Lemma 6.3 implies that $q_A(\mathbf{x}, t) \leq \kappa_d \text{Cap}(A)/R_A^d$ for $t > 0$ if $\eta_{\mathbf{x}} \geq R_A$.

The next lemma provides Harnack type estimates for q_A .

Lemma 6.4. *If $3R_A \leq |\mathbf{x}| \leq |\mathbf{y}| \leq t/R_A$, then*

$$q_A(\mathbf{y}, t) \leq \kappa_d \left[q_A(\mathbf{x}, t) + \text{Cap}(A) e^{-y/3R_A} \right] \quad (y = |\mathbf{y}|). \quad (6.21)$$

Proof. Put $R = \frac{3}{2}R_A$ and let $3R_A \leq |\mathbf{x}| \leq |\mathbf{y}| \leq t/R_A$. We use the representation

$$q_A(\mathbf{y}, t) = \int_0^t ds \int_{\partial U(R)} H_{U(R)}(\mathbf{y}, s; d\xi) q_A(\xi, t-s). \quad (6.22)$$

There exists a positive constant κ'_d such that for $s > R_A y$ ($y = |\mathbf{y}|$),

$$H_{U(R)}(\mathbf{y}, s; d\xi)/m_R(d\xi) < \kappa'_d q(y, s; R) \quad (6.23)$$

and $H_{U(R)}(\mathbf{x}, s; d\xi)/m_R(d\xi) \geq (\kappa'_d)^{-1} q(x, s; R)$, hence comparing the integral (6.22) restricted on the interval $[R_A y, t]$ with the same integral but with $\mathbf{x}$ replacing $\mathbf{y}$ we have

$$\int_{R_A y}^t ds \int_{\partial U(R)} H_{U(R)}(\mathbf{y}, s; d\xi) q_A(\xi, t-s) \leq \kappa''_d q_A(\mathbf{x}, t).$$

On the other hand

$$\begin{aligned} \int_0^{R_A y} ds \int_{\partial U(R)} H_{U(R)}(\mathbf{y}, s; d\xi) q_A(\xi, t-s) &\leq \kappa_d \text{Cap}(A) \int_0^{R_A y} q(y, s; R) ds \\ &\leq \kappa'''_d \text{Cap}(A) e^{-y/3R_A}, \end{aligned}$$

where Lemma 6.3 (ii) and Theorem 2.2 are applied for the first and second inequalities, respectively. Thus we obtain (6.21). $\square$

Lemma 6.5. *If $x/R_A > (t/R_A^2)^{1/d} > 1$, then $q_A(\mathbf{x}, t) \leq \kappa_d t^{-d/2} \text{Cap}(A)$.*

Proof. We may suppose $R_A = 1/3$. The proof is parallel to that of Lemma 6.3 but this time we consider the hitting of the ball $U(1)$ (instead of the exiting from $\mathbf{x} + U(1)$), choose $s^* \in [\frac{1}{3}t, \frac{2}{3}t]$ so that

$$q_A(\mathbf{x}_0, s^*) \leq \kappa_d t^{-1} (t \vee x^2)^{-\nu} \text{Cap}(A) \quad (6.24)$$

with any fixed $\mathbf{x}_0 \in \partial U(1)$ (which is possible owing to Lemma 6.2) and look at the decomposition

$$\begin{aligned} q_A(\mathbf{x}, t) &= \int_0^{t-s^*} ds \int_{\partial U(1)} H_{U(1)}(\mathbf{x}, s; d\xi) q_A(\xi, t-s) \\ &\quad + \int_{\mathbf{y} \notin U(1)} P_{\mathbf{x}} \left[B_{t-s^*} \in d\mathbf{y}, \sigma_{U(1)} > t-s^* \right] q_A(\mathbf{y}, s^*). \end{aligned} \quad (6.25)$$

On applying Lemma 6.4 (with $\mathbf{x}_0, s^*$ in place of $\mathbf{x}, t$) and (6.24) in turn we deduce that

$$q_A(\mathbf{y}, s^*) \leq \kappa_d [q_A(\mathbf{x}_0, s^*) + \text{Cap}(A) e^{-y/3R_A}] \leq \kappa'_d \text{Cap}(A) (t^{-d/2} + e^{-y}),$$

hence that the second term on the right-hand side of (6.25) is dominated by a constant multiple of $t^{-d/2} \text{Cap}(A)$.

Since $x^d p_s^{(d)}(x) \leq x^d p_{x^2/d}(x) \leq (d/2\pi e)^{d/2}$, it follows from Corollary 2.1 that

$$q(x, s; 1) \leq \kappa_d x^{-d} \quad \text{if } s \wedge x \geq 1, \quad (6.26)$$

which together with (6.23) shows that the inner integral of the repeated integral of the first term is at most $\kappa_d x^{-d} \int q_A(\xi, t-s) m_1(d\xi)$. Hence, on employing Lemma 6.2 as well as the assumption of the lemma the first term is dominated by $\text{Cap}(A)$ times $\kappa_d'' t^{-\nu} x^{-d} \leq \kappa t^{-d/2}$ as desired. $\square$

Lemma 6.6. *If $t > R_A^2$, then for all $\mathbf{x} \in \Omega_A$,*

$$q_A(\mathbf{x}, t) \leq \kappa_d t^{-d/2} \text{Cap}(A).$$

Proof. Suppose $R_A = 1/2$ for simplicity. Owing to the preceding lemma it suffices to show the inequality for $x \leq t^{1/d}$. The range of x may be further restricted to $x > 2R_A$. Indeed, making decomposition

$$\begin{aligned} q_A(\mathbf{x}, t) &= \int_{\mathbf{y} \in U(1)} P_{\mathbf{x}} \left[B_{t/2} \in d\mathbf{y}, \sigma_{\partial U(1)} \wedge \sigma_A \geq t/2 \right] q_A(\mathbf{y}, \tfrac{1}{2}t) \\ &\quad + \int_{\mathbf{y} \notin U(1)} P_{\mathbf{x}} \left[B_{t/2} \in d\mathbf{y}, \sigma_{\partial U(1)} \leq t/2 < \sigma_A \right] q_A(\mathbf{y}, \tfrac{1}{2}t), \end{aligned}$$

we observe that the first term on the right is $\text{Cap}(A) \times e^{-\lambda t}$ owing to Lemma 6.3 (ii) and the second is reduced to the case $x \geq 2R_A$.

Let $1 (= 2R_A) \leq x \leq t^{1/d}$ and put $r = 2t^{1/d}$. In the right-hand side of the decomposition

$$\begin{aligned} q_A(\mathbf{y}, t) &= \int_0^{t/2} ds \int_{\partial U(r)} P_{\mathbf{x}} \left[B_s \in d\xi, \sigma_{\partial U(r)} \in ds, s < \sigma_A \right] q_A(\xi, t-s) \\ &\quad + \int_{U(r) \cap \Omega_A} P_{\mathbf{x}} \left[B_{t/2} \in d\mathbf{y}, \tfrac{1}{2}t < \sigma_{\partial U(r)} \wedge \sigma_A \right] q_A(\mathbf{y}, \tfrac{1}{2}t), \end{aligned}$$

the second term is at most $\kappa_d e^{-\lambda t/r^2} \text{Cap}(A)$ for some constant $\lambda > 0$ owing to the bound $q_A(\cdot, t) \leq \kappa_d \text{Cap}(A)$ ($t \geq 1$), whereas Lemma 6.5 shows that the repeated integral is dominated by $\kappa_d t^{-d/2} \text{Cap}(A)$. Thus Lemma 6.6 has been proved. $\square$

Combined with Corollary 2.1 as well as with the last lemma the following one, virtually a corollary of Lemma 6.5, concludes the proof of proposition 6.3.

Lemma 6.7. *For each $M > 1$, there exists a constant $C_{d,M}$ such that if $2R_A \leq \sqrt{t} \leq x \leq Mt/R_A$, then*

$$q_A(\mathbf{x}, t) \leq C_{M,d} R_A^{-2\nu} \text{Cap}(A) q(x, t; R_A).$$

Proof. It follows from Corollary 2.1 that $q(x, t; a)$ is of the same order of magnitude as

$$a^{2\nu} \left(\frac{t}{ax} \wedge 1 \right)^{\frac{1}{2}(d-3)} p_t^{(d)}(x-a)$$

as $t \rightarrow \infty$ uniformly for $a > 0$ as well as for $x > 2a$. Since for $\alpha = 0$ or $\frac{1}{2}(d-3)$, the function $\varphi(s) = s^\alpha p_s^{(d)}(x)$ is increasing in the interval $0 \leq s \leq x^2/(d-2\alpha)$, this shows that for t_0 large enough,

$$q(x, s; R_A) \leq \kappa_d q(x, t; R_A) \quad \text{for } t_0 < s < t, \quad (6.27)$$

provided that $x \geq \sqrt{t} > 2R_A$. The integral (6.22) with $R = R_A$ restricted to $0 \leq s < t/2$ is at most $\kappa_d t^{-d/2} \text{Cap}(A) \int_0^{t/2} q(\mathbf{x}, s; R_A) ds$, which is negligible for the present estimate by virtue of Theorem 2.4. Applying (6.23) and (6.27) in turn, the rest of the integral (6.22) is dominated by $C'_{M,d} q(x, t; R_A) P_{m_{R_A}}[\sigma_A < t/2]$, hence by the required bound of the lemma owing to the identities (3.2) and $\text{Cap}(U(a)) = a^{2\nu}/G(1)$. $\square$

6.4 Some Upper bounds of q_A ($d = 2$)

The statements corresponding to Proposition 6.3 and Lemma 6.6 for $d = 2$ are given by the following ones.

Proposition 6.4. *There exists a universal constant C such that for all $t > R_A^2$ and $\mathbf{x} \in \Omega_A$,*

$$q_A(\mathbf{x}, t) \leq \begin{cases} \frac{C e_A(\mathbf{x})}{t(\lg(t/R_A^2))^2} & \text{for } x < \sqrt{t}, \\ C \Lambda_0(R_A x/t) p_t^{(2)}(x) & \text{for } x \geq \sqrt{t}. \end{cases}$$

Proposition 6.5. *There exists a universal constant C such that for all $t > R_A^2$ and $\mathbf{x} \in \Omega_A$,*

$$q_A(\mathbf{x}, t) \leq C \frac{e_A(\mathbf{x}) \wedge \lg(2t/R_A^2)}{t(\lg(t/R_A^2))^2}.$$

The proof of these results are similar to the corresponding ones for $d \geq 3$ but there is a crucial point to be modified. In place of Lemma 6.2 we can readily derive the following bound

$$P_{\mathbf{x}}[\sigma_A > t] \leq C e_A(\mathbf{x}) / \lg(t/R_A^2)$$

for $t > R_A^2$ by an argument analogous to that of the second half of the proof of Lemma 6.6. This bound is sharp for itself (for $x < \sqrt{t}$) but not so useful (if $m_{R_A}(e_A)$ is large) for the present purpose since its use bring an additional factor $m_{2R_A}(e_A)$. In below we use only the trivial bound $\int_{t_1}^{t_2} q_A(\mathbf{x}, s) ds \leq 1$; the same proof of Lemma 6.3 deduces from it that

$$q_A(\mathbf{x}, t) \leq C_0 \quad \text{if } t \vee \text{dist}(\mathbf{x}, A^r) \geq 1, \mathbf{x} \in \Omega_A. \quad (6.28)$$

Lemma 6.8. *There exists a universal constant C such that for $\mathbf{x} \in \Omega_A, t > 2R_A^2$,*

$$q_A(\mathbf{x}, t) \leq \begin{cases} C [t \lg(t/R_A^2)]^{-1} & \text{for } x < \sqrt{t}, \\ C q(x, t; 2R_A) & \text{for } x \geq \sqrt{t}. \end{cases} \quad (6.29)$$

Proof. Suppose $R_A = 1/2$ for simplicity. We only consider the case $x \geq 1$, to which the case $x < 1$ can be easily reduced. (See the first half of the proof of Lemma 6.6.) Let $R = 2R_A = 1$. We shall repeatedly apply the decomposition

$$q_A(\mathbf{x}, t) = I + II \quad (x > 1) \quad (6.30)$$

where

$$I = \int_0^{t/2} \int_{U(1)} H_{U(1)}(\mathbf{x}, t-s; d\xi) q_A(\xi, s) ds$$

and

$$II = \int_0^{t/2} \int_{U(1)} H_{U(1)}(\mathbf{x}, s; d\xi) q_A(\xi, t-s) ds.$$

It follows from Theorem 2.2 that for $x \geq t$, $\int_0^t q(x, s; 1) ds \leq Cq(x, t; 1)$. Substitution from it as well as from (6.28) shows $q_A(\mathbf{x}, t) \leq Cq(x, t; 1)$. For $x < t$ we apply Theorem 2.4 and the inequality $q(x, t-s; 1) \leq C'q(x, t; 1)$ ($0 < s < t/2$) to obtain that for $x > 1$,

$$I \leq Cq(x, t; 1). \quad (6.31)$$

As for II we apply the bound $q(x, s; 1) \leq C/x^2$ valid for all $s > 0$, $x > 2$. Combined with (6.31) and (6.28) this results in

$$q_A(\mathbf{x}, t) \leq C/(x^2 \vee 1) \quad (t > 2, \mathbf{x} \in \Omega_A),$$

and on using the expression

$$q_A(\mathbf{x}, t) = \int_{\Omega_A} P_{\mathbf{x}}[\sigma_A > \tfrac{1}{2}t, B_{t/2} \in d\mathbf{y}] q_A(\mathbf{y}, \tfrac{1}{2}t) \quad (6.32)$$

an easy computation deduces

$$\begin{aligned} q_A(\mathbf{x}, t) &\leq \int_{\mathbf{R}^2} p_{t/2}^{(2)}(\mathbf{y} - \mathbf{x}) \frac{C}{y^2 \vee 1} |d\mathbf{y}| \\ &\leq C' \frac{\lg t}{t} \quad (t > 2, x \geq 1). \end{aligned} \quad (6.33)$$

Now recalling the expression that defines II we obtain

$$II \leq C' P_{\mathbf{x}}[\sigma_{U(1)} < \tfrac{1}{2}t] \frac{\lg t}{t}. \quad (6.34)$$

From Theorems 2.2 and 2.3 it is readily deduced that for $x > 1, t > 2$,

$$P_{\mathbf{x}}[\sigma_{U(1)} < s] \leq \frac{C}{\lg s} \left(1 \vee \lg \frac{s}{x^2}\right) e^{-x^2/2s} e^{x/s} \quad (6.35)$$

with some universal constant C (cf. Corollary 14 of [25]), and in view of (6.31) substitution of this with $s = t/2$ into the right-hand side of the preceding inequality yields

$$q_A(\mathbf{x}, t) \leq \frac{C}{t} \left(1 \vee \lg \frac{t}{x^2}\right) e^{-x^2/2t} e^{2x/t} \quad (x > 1, t > 2), \quad (6.36)$$

Using this bound of q_A as well as (6.28) we infer from (6.32) that $q_A(\xi, t) \leq C''/t$ for $\xi \in \partial U(1)$, hence $II \leq C'''/t$, and using (6.31) again we conclude

$$q_A(\mathbf{x}, t) \leq C_2/t.$$

We have derived this bound from (6.33) and the same procedure with the bound just obtained leads to $q_A(\mathbf{x}, t) \leq C_3/t \lg t$, as desired. $\square$

Proof of Proposition 6.5. The same argument as made in the second half of the proof of Lemma 6.6 deduces Proposition 6.5 from Proposition 6.1 and Lemma 6.8. $\square$

Proof of Proposition 6.4. In view of Proposition 6.5 and the second inequality of (6.29) it suffices to show that

$$q_A(\mathbf{x}, t) \leq \kappa_d q(x, t; R_A) \quad \text{for } x \geq t/R_A.$$

(Note that here we are concerned with $\sigma_{U(R_A)}$ instead of $\sigma_{U(2R_A)}$.) For the proof we make use of a result from [27] (Theorem 2.4), which entails that there exists a family of measures $\mu_\alpha(d\xi)$, $\alpha \geq 1$, on $\partial U(a)$ such that $\mu_\alpha(\partial U(a)) < \kappa_d$ and

$$H_{U(a)}(\mathbf{x}, s; d\xi) \leq q(x, t; a) \mu_{ax/t}(d\xi) \quad \text{if } t/2 < s < t \leq ax, \quad (6.37)$$

and apply (6.30) with $U(R_A)$ in place of $U(1)$. Since $\int_0^{t/2} q_A(\xi, s) ds \leq 1$ for all $\xi \in \partial U(R_A)$ (where $q_A(\xi, \cdot)$ is regarded as the Dirac delta function if $\xi \in A^r$), (6.37) immediately gives the required bound for I in (6.30), whereas (6.34) (together with (6.35) with $s = t/2$) provides a bound of II sufficient for the present purpose. $\square$

It is noted that the procedure mentioned at the end of the proof of Lemma 6.8 may be applied once more. This no longer yields a better bound of q_A , for we must add the term I , giving rise to a term which is the same order of magnitude as $1/t \lg t$ (in view of (6.31); whereas in (6.36) we have the factor $C/t(\lg t)^2$ in place of $C/t \lg t$ and we obtain a better bound of II , which we state as a lemma.

Lemma 6.9. *With a universal constant C and $\beta_A = m_{2R_A}(e_A)$*

$$\int_0^{t/2} \int_{\partial U(R)} H_{U(R)}(\mathbf{x}, s; d\xi) q_A(\xi, t-s) ds \leq \frac{C\beta_A}{t(\lg t)^3} \left(1 \vee \lg \frac{t}{x^2}\right) e^{-x^2/t} e^{2x/t} \\ (x > R := 2R_A, t > 2).$$

Proof. This follows immediately from (6.35) and the upper bound of q_A in Proposition 6.5.

6.5 Asymptotics of the distribution of σ_A

In below we give some asymptotic estimates of the distribution function $P_{\mathbf{x}}[\sigma \leq t]$ for large time only in the case $x/t \rightarrow 0$, the case when those of the density $q_A(\mathbf{x}, t)$ are explicit enough. For the other case Propositions 6.3 and 6.4 would be enough if it is upper bound what one needs.

Proposition 6.6. *Let $d \geq 3$. Then, as $t \rightarrow \infty$ and $x/t \rightarrow 0$*

$$P_{\mathbf{x}}[t < \sigma < \infty] = \text{Cap}(A) P_{\mathbf{x}}[\sigma_A = \infty] \int_t^\infty p_s^{(d)}(x) ds (1 + o(1)) \quad (6.38)$$

uniformly for $\mathbf{x} \in \Omega_A$; and as $x \rightarrow \infty$ and $x/t \rightarrow 0$

$$P_{\mathbf{x}}[\sigma \leq t] = \text{Cap}(A) \int_0^t p_s^{(d)}(x) ds (1 + o(1)). \quad (6.39)$$

The proposition above follows from Theorem 3.1: (6.38) is immediate, whereas for the proof (6.39) only the case $\lim(\lg x)/\lg t = 1$ needs to be taken care of: in that case one may apply Propositions 6.3 to see that $p_{\mathbf{x}}[\sigma_A < t/2]$ is negligible. It is noted that

$$\int_0^t p_s^{(d)}(x) ds = (2\pi)^{-d/2} 2^\nu x^{-2\nu} \int_{x^2/2t}^\infty e^{-y} y^{\nu-1} dy$$

and similarly for $\int_0^t p_s^{(d)}(x) ds$.

Proposition 6.7. *Let $d = 2$. Then, uniformly for $\mathbf{x} \in \Omega_A$, as $t \rightarrow \infty$ and $x/t \rightarrow 0$*

$$\begin{aligned} \text{(i)} \quad P_{\mathbf{x}}[\sigma_A > t] &= \frac{2e_A(\mathbf{x})}{\lg t} \left(1 + O\left(\frac{1}{\lg t}\right) \right) \quad \text{for } x \leq \sqrt{t}, \\ \text{(ii)} \quad P_{\mathbf{x}}[\sigma_A \leq t] &= \frac{1}{2\pi \lg(t/x)} \int_{x^2/2t}^{\infty} e^{-y} y^{-1} dy (1 + o(1)) \quad \text{for } x \geq \sqrt{t/\lg t}. \end{aligned}$$

Proof. For (i) use $\int_t^{\infty} (1 - e^{-x^2/2s}) [s(\lg s)^2]^{-1} ds \leq Cx^2/t(\lg t)^2$. □

Acknowledgments. The author wishes to thank the anonymous referee for his carefully reading the original manuscript and pointing out the non-trivial mistakes and incorrect arguments perpetrated therein.

References

- [1] R. F. Bass, Probabilistic techniques in analysis, Springer, 1995
- [2] M. van den Berg, Heat flow, Brownian motion and Newtonian capacity, Ann. Inst. H. Poincaré -PR. **43** (2007), 193-214.
- [3] M. van den Berg and E. Bolthausen, On the expected volume of the Wiener sausage for a Brownian bridge, Math. Z. **224** (1997), 33-48.
- [4] R. M. Blumenthal and R. K. Gettoor, Markov processes and potential theory, Academic Press, 1968
- [5] T. Byczkowski, J. Malecki and M. Ryznar, Hitting times of Bessel processes, Potential Anal. **38** (2013), 753-786.
- [6] H.S. Carslaw and J.C. Jaeger, Conduction of heat in solids, Clarrendon press, Oxford, 1992.
- [7] K. J. Falconer, The geometry of fractal sets, Cambridge Univ. Press, 1985
- [8] Y. Hamana and H. Matumoto, The probability distributions of the first hitting times of Bessel processes, to appear in TAMS., available at arXiv:1106.6132.
- [9] Y. Hamana and H. Matumoto, The probability densities of the first hitting times of Bessel processes, J. Math-for-Ind. **4B** (2012), 91-95.
- [10] G. Hunt, Some theorems concerning Brownian motion, TAMS. **s1** (1956), 294-319.
- [11] G. Hunt, Markov processes and potentials I, II and III, Ill. J. Math. **1** (1957), 44-93, 316-369, **2**(1958) 151-213.
- [12] K. Itô and H.P. McKean, Jr, Diffusion processes and their sample paths. Springer, 1965
- [13] Joffe, A, Sojourn times for stable processes, Thesis, Cornell U., 1959
- [14] O. D. Kellog, Foundations of potential theory, Springer, 1929
- [15] I. McGillivray, Large time volume of the pinned Wiener sausage, J. Funct. Anal., **170** (2000), 107-140.
- [16] I. McGillivray, The spectral shift function for planar obstacle scattering at low energy, Math. Nachr. **286** (2013), 1208-1239.

- [17] S. C. Port and C. J. Stone, Brownian motion and classical potential theory, Academic press, (1978)
- [18] F. Spitzer, Some theorems concerning 2-dimensional Brownian motion, TAMS. **87** (1958), 187-197.
- [19] F. Spitzer, Electrostatic capacity, heat flow and Brownian motion, ZW. **3** (1964), 187-197.
- [20] D. W. Stroock, Probability, 2nd ed., Cambridge Univ. Press, 2011
- [21] E.C. Titchmarsh, Eigenfunction expansions: Part one, 2nd ed., Oxford Univ. Press, 1962
- [22] K. Uchiyama, The mean number of sites visited by a pinned random walk, Math. Zeit. **261** (2009), 277-295.
- [23] K. Uchiyama, Asymptotic estimates of the distribution of Brownian hitting time of a disc., J. Theor. Probab., **25** (2012), 450-463. / Erratum, J. Theor. Probab. **25** (2012), issue 3, 910-911 .
- [24] K. Uchiyama, The expected area of Wiener sausage swept by a disk. Stoch. Proc. Appl. **123** (2013), 191-211.
- [25] K. Uchiyama, Asymptotics of the densities of the first passage time distributions of Bessel diffusion, Trans. Amer. Math. Soc. **367** (2015), 2719-2742.
- [26] K. Uchiyama, The expected volume of Wiener sausage for a Brownian bridge joining the origin to a point outside a parabolic region, RIMS Kokyuroku 1855, Probability Symposium (2013), 10-18.
- [27] K. Uchiyama, Density of space-time distribution of Brownian first hitting of a disc and a ball, available at: <http://arxiv.org/abs/1404.4745v3>
- [28] K. Uchiyama, The mean number of sites visited by a random walk pinned at a distant point, Electron. Commun. Probab. **20** (2015), no.17, 1-9.
- [29] G. N. Watson, A treatise on the theory of Bessel functions, 2nd ed., Reprinted by Springer, 1995
- [30] N. A. Watson, Introduction to heat potential theory, Mathematical surveys and monographs, **182** AMS. (2012).